

Full length article



Towards a marine strategy for the deep Mediterranean Sea: Analysis of current ecological status

R. Danovaro^{a,b,*,1}, E. Fanelli^{a,1}, M. Canals^{c,1}, T. Ciuffardi^{d,1}, M.-C. Fabri^{e,1}, M. Taviani^{b,f,g,1}, M. Argyrou^h, E. Azzurro^{b,i}, S. Bianchelli^a, A. Cantafaro^j, L. Carugati^a, C. Corinaldesi^k, W. P. de Haan^c, A. Dell'Anno^a, J. Evans^j, F. Fogliani^f, B. Galil^l, M. Gianni^m, M. Goren^l, S. Greco^b, J. Grimaltⁿ, Q. Güell-Bujons^c, A. Jadaud^o, L. Knittweis^j, J.L. Lopezⁿ, A. Sanchez-Vidal^c, P. J. Schembri^j, P. Snelgrove^p, S. Vaz^o, the IDEM Consortium

^a Department of Life and Environmental Sciences, Polytechnic University of Marche, 60131, Ancona, Italy

^b Stazione Zoologica Anton Dohrn Naples, 80122, Naples, Italy

^c CRG Marine Geosciences, Department of Earth and Ocean Dynamics, Faculty of Earth Sciences, University of Barcelona, 08028, Barcelona, Spain

^d Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Department for Sustainability, S. Teresa Marine Environment Research Centre, 19100, La Spezia, Italy

^e Institut Français de Recherche pour l'exploitation de la Mer (Ifremer), Département Océanographie et Dynamique des Ecosystèmes, 83500, La Seyne sur Mer, France

^f Institute of Marine Sciences (ISMAR), CNR, 40129, Bologna, Italy

^g Biology Department, Woods Hole Oceanographic Institution, MA, 02543, USA

^h Department of Fisheries and Marine Research (DFMR), 1416, Nicosia, Cyprus

ⁱ Institute for Environmental Protection and Research (ISPRA) STS Livorno, 57122, Italy

^j Department of Biology, University of Malta, Msida, MSD2080, Malta

^k Department of Sciences and Engineering of the Materials, Environment and Urban Planning, Polytechnic University of Marche, Italy

^l The Steinhardt Museum of Natural History, Israel National Center for Biodiversity Studies, Tel Aviv University, Tel Aviv, 69978, Israel

^m Deep Sea Conservation Coalition, Postbus, 59681, Amsterdam, Netherlands

ⁿ Department of Environmental Chemistry, Institute of Environmental Assessment and Water Research (CSIC), 08034, Barcelona, Spain

^o UMR Marbec, Ifremer, IRD, Université de Montpellier, CNRS, 34203, Sète Cedex, France

^p Departments of Ocean Sciences and Biology, Memorial University of Newfoundland, St. John's, NL A1C 5S7 Canada

ARTICLE INFO

Keywords:

Marine strategy framework directive
Deep-sea ecosystems
Mediterranean basin

ABSTRACT

The Marine Strategy Framework Directive (MSFD), introduced in June 2008, was adopted to achieve a Good Environmental Status (GES) in the EU's marine waters and to protect resources of socio-economic interest. The MSFD exerts to the marine area over which a Member State exercises jurisdictional rights in accordance with the United Nations Convention on the Law of the Sea (UNCLOS), including the deep-sea waters, seafloor and sub-seafloor of the Exclusive Economic Zones (EEZ). However, currently the MSFD focuses on coastal habitats and the shallow-water seafloor to the detriment of the deeper habitats. Despite the huge dimension of the deep sea (below 200 m of depth) covering more than 65% of the Earth's surface and including >95% of the global biosphere, the relevance of the dark portion of the seas and oceans is still almost completely neglected. Given the important bi-directional links between shallow and deep ecosystems, there is a clear need for extending the implementation of the MSFD into the deep sea, to define a sound ecosystem-based approach for the management and protection of deep-sea ecosystems and attain GES. We assembled data on drivers, anthropogenic pressures and impacts concerning the MSFD descriptors pertaining to the Mediterranean deep sea. We list deep-sea monitoring activities and the main sources providing benchmark conditions, and discuss knowledge and geographic coverage gaps. MSFD descriptors apply to the deep sea as to coastal waters, and ought to be monitored contemporaneously. We provide recommendations for guidelines for future deep-sea monitoring in the Mediterranean Sea.

* Corresponding author. Department of Life and Environmental Sciences, Polytechnic University of Marche, 60131, Ancona, Italy.

E-mail address: r.danovaro@univpm.it (R. Danovaro).

¹ These authors have equally contributed to the manuscript.

<https://doi.org/10.1016/j.marpol.2019.103781>

Received 24 July 2019; Received in revised form 31 October 2019; Accepted 8 December 2019

Available online 27 December 2019

0308-597X/© 2019 Published by Elsevier Ltd.

1. Introduction

1.1. The Mediterranean Sea

The Mediterranean is a semi-enclosed basin between the European and African coasts with the narrow and shallow Strait of Gibraltar connecting its waters and life to the Atlantic Ocean. The Suez Canal creates a man-made connection to the Red Sea, recently doubled, which allows the penetration of tropical Indo-pacific species [1]. Concomi-

seamounts, continental rise deposits, mud volcanoes and extreme environments such as hydrothermal vents, cold seeps and deep-hypersaline anoxic basins [3,11–14]. Even seemingly “featureless” soft bottom habitats host unique and vulnerable species and habitats (e.g. sponge fields, gorgonian and pennatulacean meadows) [3].

The Mediterranean basin is threatened by multiple stressors associated with the rapid expansion of coastal populations, urbanization, changes in agricultural, industrial and shipping patterns, overfishing and exploration and extraction of offshore minerals and hydrocarbons,

List of acronyms and abbreviations

ABNJ	Areas Beyond National Jurisdiction	LIW	Levantine Intermediate Water
ACCOBAMS	Agreement on the Conservation of Cetaceans in the Black Sea Mediterranean Sea and Contiguous Atlantic Area	MBES	Multibeam Echosounder
AUV	Autonomous Underwater Vehicle	MEDIAS	Mediterranean International Acoustic Survey
CBD	Convention on Biological Diversity	MEDITS	Mediterranean International Trawl Survey
CFP	Common Fisheries Policy	MS	Member States
CWC	Cold-Water Corals	MSFD	Marine Strategy Framework Directive
DCF	Data Collection Framework	MSY	Maximum Sustainable Yield
DSF	Deep-Sea Fisheries	NIS	Non-Indigenous Species
DSWC	Dense Shelf Water Cascading	OMZ	Oxygen Minimum Zones
EEZ	Exclusive Economic Zone	PAH	Polycyclic Aromatic Hydrocarbons
EFH	Essential Fish Habitats	POM	Particulate Organic Matter
EIA	Environmental Impact Assessment	POP	Persistent Organic Pollutants
EMT	Eastern Mediterranean Transient	ROV	Remotely Operated Vehicle
EwE	Ecopath with Ecosim	SAC	Scientific Advisory Committee
FAO	Food and Agricultural Organization	SCA	Stomach Content Analysis
FRA	Fishery Restricted Areas	SSB	Spawning Stock Biomass
GFCM	General Fisheries Commission for the Mediterranean	SIA	Stable Isotope Analysis
GSA	Geographical Sub Area	SSS	Side-Scan Sonar
ICCAT	International Commission for the Conservation of Atlantic Tunas	STECF	Scientific, Technical and Economic Committee for Fisheries
		VME	Vulnerable Marine Ecosystem
		UNCLOS	United Nations Convention on the Law of the Sea
		UNGA	United Nations General Assembly

tantly, the Mediterranean Basin is experiencing major climatic-related changes that are strongly influencing its oceanography in terms of water mass characteristics (e.g., temperature, salinity, dissolved oxygen), currents, nutrients and relative sea levels. The interplay of these factors has, since historical times, strongly influenced the diversity and colonization of the Mediterranean Sea [2,3].

Unique hydrology characterizes the present-day Mediterranean, including microtidal regime, oligotrophy, high salinity (37.5–39.5 psu), homoeothermic temperatures from 300 to 500 m to the bottom, with values at the seafloor ranging from ca 13–13.5 °C in the western basin to 13.5–15.5 °C in the eastern basin, and the almost complete lack of thermal boundaries [4]. These features uniquely make the Mediterranean one of the “warmest” deep-sea basins of the world. Strong energy gradients also characterize the Mediterranean, with primary production and food supply to the deep decreasing from the western to the eastern region of the basin and from shallower to deeper waters [5].

These historical, topographic and environmental characteristics complicate deep-sea biodiversity patterns of the Mediterranean Sea but raise intriguing questions. Numerous studies document that the Mediterranean Sea, although modest in size (0.82% and 0.32% of the global ocean surface and volume, respectively; [6], is a biodiversity hot spot with overall ca 17000 species, which represent 7.5% of the species richness of the oceans [7–9]. However, although data on the species richness of its deeper habitats are incomplete (two thirds of the deep species – excluding prokaryotes – have not been censused yet; [3,8,10], it appears that the biodiversity of the deep Mediterranean basin is lower than that of other oceans [3].

The biodiversity of the deep Mediterranean Sea depends largely from the heterogeneity of habitats, which include submarine canyons and

which exert increasing pressures through habitat destruction, chemical pollution, and dumping of waste and litter [15,16]. In concert with climate change, these stressors may act synergistically to affect the dynamics, and potentially the resilience, of fragile deep ecosystems [17, 18]. Direct stressors and processes also occur on the adjacent shelf and in the epi-mesopelagic zones, including Dense Shelf Water Cascading (DSWC) events down the continental slope, open-sea convection and severe coastal storms, which may transport sediments and organic matter to the continental slope and beyond, influencing deep-sea biodiversity and ecosystem functioning [19–23]. In particular, bottom-contacting fisheries, specifically bottom-trawling and longlines, represent the most significant anthropogenic threats to deep-water biota, severely impacting sensitive habitats and species such as cold-water corals (CWCs) and/or sponge gardens [24,25]. Additional evidence attributes a significant proportion of deep-sea litter to the fishing industry [26–29], along with land- and ship-based sources [30].

Given the increasing pressures on deep-sea habitats, scientists and managers are becoming conscious of the need to develop standardized tools and harmonized observation systems for long-term biological monitoring, in order to enable the collection of scientifically-validated data and a better understanding of the consequences of the present and future anthropogenic impacts [31–33].

1.2. Implementing the marine strategy framework directive in the deep Mediterranean Sea

The Marine Strategy Framework Directive (MSFD 2008/56/EC) represents the EU’s Integrated Maritime Policy tool to achieve Good Environmental Status (GES) of marine waters, with an initial target for

2020. The MSFD applies to the area of marine waters over which a Member State exercises jurisdictional rights in accordance with the UNCLOS (see Fig. 1, for the definition of territorial waters and EEZs in the Mediterranean). These include also deep-sea waters, seabed and sub-seafloor. At present, MSFD implementation focuses mostly on coastal habitats or those impacted by commercial fisheries [34]. However, over long-time scales, global nutrient and carbon cycles depend on a functioning deep sea (e.g. Ref. [35]). Moreover, the life-cycle stages of some coastal species use offshore environments, thus achieving GES for marine ecosystems associated with continental shelves, must link to the achievement of GES for deep Mediterranean environments and Areas Beyond National Jurisdiction (ABNJ or “High Seas”). Otherwise, the MSFD will largely disregard the precautionary principle and undermine an ecosystem-based approach to marine management.

An effective governance and management of the Mediterranean Sea requires consideration of the complexity of these environmental issues, and meaningful international cooperation [36]. Given the trans-boundary nature of most of the deep waters, their inclusion in MSFD complicates the requirement for each Member State to apply the Directive to areas within its national jurisdiction. This emphasizes the need for Member States (MS) to cooperate in order to ensure coordinated and harmonized development of marine strategies at the scale of region/sub-region in the Mediterranean Basin, where EU MS and developing countries co-exist.

MSFD implementation currently suffers from a lack of standardized and consistent methodology for deep waters. To address this gap, we identify approaches, variables, and methodologies to enable MSFD implementation in the deep Mediterranean Sea. This synthesis summarises available information on MSFD descriptors for the deep

Mediterranean Sea, with respect to the criteria listed in the European Commission Decision (COMM/DEC/2017/848), and anthropogenic pressures, uses and human activities affecting the marine environment (Table 2 of Annex III of COMM/DEC/2017/848).


2. State of the knowledge of MSFD descriptors in the deep mediterranean

2.1. Descriptor 1: biological diversity

Descriptor 1 (D1) states that “The quality and occurrence of habitats and the distribution and abundance of species are in line with prevailing physiographic, geographic and climatic conditions” (MSFD, 2008/56/EC, Annex I, summarised in Table 1). The species groups specified in Part II of the Annex to COMM/DEC/2017/848, these include birds, mammals, reptiles, fish and cephalopods, some of which are present, diverse and abundant in the deep sea, such as fishes and cephalopods, in addition to deep diving and feeding cetaceans. Deep-sea organisms play an important role in marine food webs, either as predators or as important prey of a large set of high trophic level predators, including other fishes and cephalopods and marine mammals [37–39].

Data on these components from the deep Mediterranean Sea are largely included in the MEDITS database, which also represent the only extensive time series available for the deep Mediterranean [40], although with the limitation that it is a destructive sampling method with discrete sampling time, which exclude the possibility to detect any displacements of demersal species [41,42]. MEDITS is funded as part of the EU Data Collection multi-annual sampling program (DC-MAP), which limits the sampling frequency to yearly surveys confined to the

Jurisdictional Continental Shelf and Deep Mediterranean Sea



Approximative percentages of Jurisdictional Continental Shelf per Country related to the total area of deep sea, calculated with the 200 meters isobath limit, in the Mediterranean Sea: Albania (AL): 0,3%; Algeria (DZ): 6,1%; Croatia (HR): 0,5%; Cyprus (CY): 4,8%; Egypt (EG): 7%; France (FR): 3,5%; Greece (EL): 20,6%; Israel (IL): 1,1%; Italy (IT): 21,2%; Lebanon (LB): 0,95%; Lybia (LY): 15,1%; Malta (MT): 2,3%; Montenegro (ME): 0,15%; Morocco (MA): 0,6%; Spain (ES): 11,1%; Syria (SY): 0,5%; Tunisia (TN): 1,5%; Turkey (TR): 2,5%.

Fig. 1. Jurisdictional Continental Shelf and deep Mediterranean Sea, with indication of jurisdictional continental shelf per each country (including non-EU countries).

Table 1

List of the descriptors, their definition if GES is achieved and if they are a state or pressure descriptors. Only D3 is both a state and pressure descriptor as it related to aspects such as the level of fishing activity (pressure) and population age, size distribution and biomass indices (state).

Descriptor	GES	State	Pressure
1	Biodiversity is maintained	X	
2	Non-indigenous species do not adversely alter the ecosystem		X
3	The population of commercial fish species is healthy	X	X
4	Elements of food webs ensure long-term abundance and reproduction	X	
5	Eutrophication is minimised		X
6	The sea floor integrity ensures functioning of the ecosystem	X	
7	Permanent alteration of hydrographical conditions does not adversely affect the ecosystem		X
8	Concentrations of contaminants give no effects		X
9	Contaminants in seafood are below safe levels		X
10	Marine litter does not cause harm		X
11	Introduction of energy (including underwater noise) does not adversely affect the ecosystem		X

northern part of the Mediterranean Basin. MEDITS mostly targets demersal fish (including deep-sea sharks), but includes also commercial invertebrates and other macro and mega-invertebrates (as by-catch species). MEDITS provides detailed information on their abundance and biomass, including the population structure of several species (including length frequency distributions by sex and maturity stages for different target species), which allow us to obtain information on the size spectra, maturity ogives, sex ratios and mortality rates. This information contributes to both the census of shallow and deep marine biodiversity and stock assessments carried out by the GFCM and the STECF of the European Commission (see Refs. [43,44] [see Descriptor 3 below]. In the case of meso- and bathypelagic species [45]; i.e. [38, 46–49], species of non-commercial interest, hard bottom habitats between 200 and 800 m depth, and in general all habitats below 800 m depth, only scattered information without temporal datasets exist.

Another important gap concerns the smaller size biota, such as meiofauna, which are a key component in the deep-sea ecosystems and are driven by water depth, regional setting and geomorphological characteristics of the deep Mediterranean habitats [50]. Meiofauna are highly diversified (possibly hyper-diverse), and play a fundamental ecological role in the biogeochemical cycles and in food webs and are sensitive to environmental and anthropogenic pressures [51]. Since this component, which increases its ecological relevance, in terms of abundance and functional role, with increasing water depths [52], has been recently suggested for inclusion in the D1 for the implementation of the MSFD [53,54,287]. It is even more evident that it should be taken into consideration in the implementation of the MSFD in the deep sea.

The MSFD deep-sea habitats included in the COMM/DEC/2017/848 are: a) upper bathyal rocks and biogenic reefs, b) upper bathyal sediments, c) lower bathyal rock and biogenic reef, d) lower bathyal sediments, and e) abyssal seafloor. These include other benthic habitats such as: canyons (which may include rocky and sedimentary substrates), rocky bottoms with coral banks (including Cold-water corals) or large bivalves, different types of sedimentary bottoms in bathyal or abyssal plains (muds, sands or coarser sediment), chemosynthetic ecosystems (hydrothermal vents and mud volcanoes), and seamounts.

Previous studies on the deep Mediterranean Sea reported a west–east decreasing gradient of food availability [5,55], which explains the presence of a significant decreasing gradient in the abundance and biomass of most deep-sea benthic components along that gradient, from meiofauna to megafauna [50,56,57]. The CWCs apparently follow the same gradient [58,59], and the presence of Levantine Intermediate Water (LIW) likely strongly influences their distribution [23,58,60].

Trawl surveys provide most of the available information on deep-sea habitats and their characteristics (see Table 1, Annex III, MSFD), but only for soft bottom habitats including those dominated by *Isidella elongata* and *Funiculina quadrangularis* [61,62]. Oceanographic cruises using ROVs offer the possibility to conduct non-destructive image and sample collections able to contribute significantly to the study of deep-sea habitats. Most ROV surveys to date have focused on CWC habitats and coral gardens, and provide important information on the composition, abundance, and biomass of the communities within these habitats [63,64,59,65–72,288]. Most available information focuses on deep-sea canyons [73], seamounts [74] and mud volcanoes [75], with major data gaps for deep-sea pelagic habitats, notwithstanding there is an increasing information on deep-water zooplankton and micronekton [31,76–82]. Descriptor 1 is directly linked to D2, D3, D4 and D6 (habitats), and monitoring efforts in the deep sea can therefore gather contextual information on all these descriptors. The ecosystem criteria listed in COMM/DEC/2017/848, which link Descriptors 1 and 4, consider trophic guilds. These are highly relevant to the deep sea and can therefore be immediately described, as the already available data would provide the required background information.

2.2. Descriptor 2: non-indigenous species introduced by human activities

The number of recorded Non-Indigenous Species (NIS) in the Mediterranean Sea greatly exceeds that in other European seas [83,84]. Their establishment alters biotic assemblages and ecosystem functions [48, 85–90]. The Suez Canal is an important pathway for Red Sea species, which indeed represent 2/3 of the NIS in the Mediterranean Sea [88]. In the past, it was assumed that NIS could establish only in shallow waters, however, the deep sea is not immune to species invasions. NIS have been rarely documented in the deep sea, a notable exception is the red king crab *Paralithodes camtschaticus* in the Barents Sea (Jørgensen and Nilsen, 2011). Yet, a growing number of Erythraean species were reported from the deeper part of the continental shelf, beyond the shelf break and in the upper slope [91–94]. For example, the lethally poisonous silver-cheeked toadfish, *Lagocephalus scleratus*, has been collected from 350 to 400 m depth off Spain [289]. The invasive lionfish, *Pterois miles*, that was initially present only in the upper shelf has been recently recorded at depths down to 110–150 m [95,96]. In the southern Levantine Sea, three carnivorous species of Erythraean origin have been observed at 200 m depth: the crocodile toothfish *Champsodon nudivittis*, the burrowing goby *Trypauchen vagina* and the red-eye round herring *Etrumeus golanii* [97]. The presence of deeper dwelling populations suggests that thermal niche assessments based only on a species' native range may underestimate their ability to tolerate lower temperatures [98]. Wider thermal tolerance of some Erythraean species may facilitate their bathymetric and geographic expansion to depths where unique, diverse, and fragile mesophotic 'animal forests' occur. The lately observed "descent" of NIS from the upper to lower continental shelf may be an indication of temperature-dependent range expansion at increasing water depths, and appears to be accelerating. Therefore, even if abundances of NIS at levels of true invasions have not been reported yet in the deep Mediterranean, these vulnerable environments should be monitored also for D2, as they could be future targets of NIS invasions.

2.3. Descriptor 3: populations of commercially exploited fish and shellfish

Descriptor 3 determines that Member States should maintain commercially exploited stocks of fish and shellfish in a healthy state. This descriptor implies sustainable exploitation that does not exceed the Maximum Sustainable Yield (MSY), i.e. the maximum yield catch that can be taken annually without reducing the fish stock productivity. Heavy fishing pressures, such as overexploitation or overfishing, produce negative environmental and socio-economic impacts, ranging from loss of significant potential yield of targeted stocks to severe stock depletion and fisheries collapse [99]. Overfishing can also reduce fish

stocks dramatically to the point where they lose internal genetic diversity and, with it, their capacity to adapt to environmental change [100,101]. Fish communities may also change, such as altered size structures, when fisheries target or discard particular-sized individuals of a species, may potentially affect predator and prey dynamics [102], i. e. Descriptor 4 addresses the question of trophic relationships and marine food webs. The MSFD builds on existing EU legislation such as the Common Fishery Policy (CFP), and the criteria describing stock status follow internationally acknowledged best practices. The SAC of the GFCM, the STECF of the European Commission and the ICCAT (for highly migratory species, such as tunas or swordfishes, which account for more than 10% of the value of the total catches in the Mediterranean) collectively monitor exploitation of fisheries resources in the Mediterranean marine sub-regions. The FAO and GFCM oversee collection of fisheries monitoring data in the Mediterranean Sea within GSAs (Geographical Sub Areas management divisions, according to resolution GFCM/33/2009/2, www.gfcm.org, for the correspondence between GSA numbers and their names Fig. 2), often assessing stocks over one or several GSAs. However, the MSFD sub-regions do not match with the GSAs. Furthermore, when we focus our attention on depths >200 m, the distinction between shallow and deep-water species is often irrelevant because distribution, exploitation and assessment of many stocks often cover wide depth ranges.

Descriptor 3 stipulates the need for fishery-induced mortality, yielding (but not exceeding) MSY (D3C1), and that populations of all commercially exploited species should remain within safe biological limits (D3C2), with a population age and size distribution (D3C3) indicative of a healthy stock. Fulfilling D3 criteria for deep-sea stocks

requires: (1) sustainable exploitation consistent with high long-term yields, (2) maintaining full reproductive capacity in order to maintain stock biomass, and (3) maintaining or increasing the proportion of older and larger fish/shellfish, an indicator of a healthy stock. Achieving GES also for a deep-sea stock requires fulfilling all three of these attributes and, for the reasons highlighted above, D3 indicators require trans-national cooperation at the level of each MSFD sub-region.

In the Mediterranean Sea, the enforcement of the CFP and, more recently, of the MSFD, continues to fall far short of achieving its objectives for exploited living marine resources (e.g., Refs. [43,103]. Notwithstanding the enforcement of the EU-Data Collection Regulation [290] in the early 2000s by all EU Member States, and the rapid increase in the number of assessed stocks by the GFCM and the STECF, industries continue to exploit Mediterranean Sea marine resources above MSY levels, with few signs of population recovery [43,44,103].

Management practices of DSF and VMEs in the Mediterranean were reviewed in 2016 and 2017 [104,105]. UNGA Resolutions 51/2006 and the FAO International Guidelines for the Management of Deep-sea Fisheries in the High Seas [106], identify DSF in the Mediterranean Sea, as those: a) using bottom-contact gears or b) using deep-pelagic trawls and c) targeting species associated with the sea floor between 300/400 m and 1000 m depth [104,105]. This grouping considers shallower fisheries that extend below 400 m. These reports identify deep-water red shrimps (*Aristaeomorpha foliacea* and *Aristeus antennatus*) as the primary DSF targets in the Mediterranean deep-sea habitats, which are harvested mostly at 300/400–800 m depths, along with European hake (*Merluccius merluccius*), Norway lobster (*Nephrops norvegicus*) and deep-water rose shrimp (*Parapenaeus longirostris*). In




Fig. 2. Location of GFCM Geographical Sub-Areas (GSAs) and MSFD Mediterranean national sub-regions. FAO divisions are shown in thumbnail map.

addition, gillnet fisheries and demersal long-liners target both *M. merluccius* and the blackspot seabream (*Pagellus bogaraveo*). Also in Spain, deep-sea fisheries target *Plesionika* shrimps below 300 m depth (see also IDEM 2018a). GFCM banned the use of towed dredges and trawl nets at depths beyond 1000 m in 2005 (Recommendation GFCM/29/2005/1), protecting over ca. 1,700,000 km² of Mediterranean Sea seafloor habitats (about 59% of the GFCM area of application) [107].

To date, the GFCM has established a number of Fishery Restricted Areas (FRAs) to protect Essential Fish Habitats (EFHs) or/and VMEs from excessive fishing mortality or the significant adverse impact of fishing activities through bottom-contact fishing gears, respectively. The FRAs encompass a total marine area of ca. 22,500 km² [107]. Four of the FRAs were declared within a multiannual management plan for deep-sea fisheries, in order to protect EFHs for spawners of several species that are heavily exploited, to maintain habitat of the continental slope (canyons and submarine canyons), and to preserve all the species of the area (commercially exploited or not), i.e., one in the Gulf of Lion in France [108], one in the Jabuka-Pomo Pit [109] and two FRAs south of Sicily. The existence of management plans, however, does not necessarily imply regular completion of accurate stock evaluation. Moreover, the existence of the FRA does not necessarily imply banning bottom trawling and may simply represent an effort management tool (to prevent further effort increase as seen in freezing fishing effort in the Gulf of Lion FRA).

Member States shall establish a list of commercially exploited species to which the criteria apply in each assessment area through regional or sub-regional cooperation, and update that list for each six-year evaluation period, taking into account Council Regulations (EU) 1251/2016, 1380/2013, 1343/2011, and 1967/2006, in accordance with article 43 (3) of the Treaty on the Functioning of the European Union, article 9 of Regulation (EU) No 1380/2013 and article 19 of Regulation (EC) No 1967/2006.

The MSFD criteria available for coastal environments cannot often be directly utilised for deep-sea species. This is because stock assessments are limited and not sufficiently monitored, hampering the assessment of stock exploitation at MSY (Criterion D3C1). The available information decreases eastwards and southwards. In addition, a gap of knowledge is also present in terms of time-series coverage of Spawning Stock Biomass (SSB) (Criterion D3C2) trend data hampering the possibility to define appropriate reference points.

The third criterion, i.e. “Healthy age and size structure” (criterion D3C3), assumes that a stock with sufficient large, and therefore old, fish corresponds to a healthy stock, thus reflecting good status. The larger and older fish stocks indicate healthier conditions, but this criterion has not been developed because GES lacks accepted thresholds [15].

This gap suggests a need to identify and test suitable indicators, metrics, and thresholds for populations and age size distributions for each deep-sea stock [110,111]. In conclusion, our analysis points out that there is a potential to inform D3 criteria, but more data need to be collected in the future to propose sound stocks analyses and reference conditions, likely through the extension of the EU Data Collection Multiannual Programme (DC-MAP, EU Regulation 2016/1701) to include more deep-sea species.

2.4. Descriptor 4: marine food webs

Descriptor 4 addresses the functional aspects of marine food webs, especially the rates of energy transfer within the system and levels of productivity in key components. In the context of the MSFD, this descriptor reaches GES when “All elements of the marine food webs, to the extent that they are known, occur at normal abundance and diversity and population levels capable of ensuring the long-term abundance of the species and the retention of their full reproductive capacity” (MSFD, 2008/56/EC, Annex I).

Deep-sea food webs critically depend on input of organic carbon from the photic zone [112]. The microbial loop and viral infections can

play an important role in the functioning of deep-sea food webs both in the water column and in sediments, and in the control of the biogeochemical cycles [55]. At the same time, deep-sea meiofauna represents a potential basic linkage in energy transfer from the benthic detritus and microbes to the macro-megafauna and demersal fishes [113,114].

D4 is one of the most controversial MSFD Descriptors in terms of protocols, criteria, and thresholds [115]. D4 is generally investigated along with Descriptors D1 and D6 or D3. Studies of food web properties typically utilize two different approaches: a) Stomach Contents’ Analyses (SCA) and b) Stable Isotope Analyses (SIA), and fatty acid trophic markers. Modelling techniques, in contrast, can provide insights regarding the potential structure of food webs [116]. SIA identified three trophic levels among deep-sea supra-benthos [117] and four levels within macrozoobenthos [77,118] and macrozooplankton/micronekton [78]. Fishes, decapods, and cephalopods dominate higher trophic levels in deep-sea demersal communities. Despite the availability of a large dataset for the deep Mediterranean Sea [119], this dataset includes few commercial species from bathyal depths: the European hake *Merluccius merluccius* and the greater forkbeard *Phycis blennoides*, some sharks (mostly *Etmopterus spinax* and *Galeus melastomus*), decapods (the red shrimps *Aristaeus antennatus* and *Aristaeomorpha foliacea*, the rose shrimp *Parapenaeus longirostris* and the Norway lobster *Nephrops norvegicus*). We lack sufficient data on other dominant deep-sea species, such as macrourids, or key predators such as deep-sea sharks (e.g., *Centroscyms coelolepis*; [120–122]).

Most studies on trophic functional groups have focused on macro- and megafauna (i.e. fish, decapods, cephalopods and echinoderms), whereas few studies are available on other biotic compartments such as meiofauna or mesozooplankton [3]. The northwestern Mediterranean portion is better studied [38,47,48,77,78,117,123,124], whereas the Ionian and Aegean seas have been much less investigated [76,125–128]. On the other hand, these types of studies rarely consider some key species/taxa, such as mesopelagic fishes or megazooplankton [49,79].

COMM/DEC/848/2017 sets criteria and methodological standards for monitoring and assessment of GES within the theme “Ecosystems”. For example, selection criteria require that at least one of the three trophic guilds monitored should focus on primary producers. This criterion is the major drawback for this descriptor given that, aside from the few, very localized ecosystems that depend on chemosynthesis (hydrothermal vents, cold seeps, or wood and whale falls; [129,130]), the vast majority of the deep sea lacks primary production. The data gap between experimental and functional data adds further complication.

Stable isotope analysis may comply with the primary criterion D4C1 (diversity of trophic guilds) and the secondary criterion D4C3 (distribution of individuals across the trophic guild). Moreover, in combination with abundance, biomass, and other biological data available from MEDITS data, it may offer inputs into ecosystem models that could generate useful outputs, such as identification of unrecognized keystone species, a gap not presently considered. Italy addresses D4 under its fishery monitoring program (i.e. D3) and specifically with three sub-programs aimed at: i) defining, testing, and applying ecosystem indicators through models (essentially EwE, <http://www.ecopath.org>); ii) identifying functional groups through the application of stable isotope analysis of monitored species within the DCF; and iii) integrating analysis of commercial species with those for benthos, zooplankton and Particulate Organic Matter (POM) samples every three years. Spain and France recently introduced SIA and SCA of species collected during MEDITS or MEDIAS surveys for use in D4.

In conclusion, the analysis of D4 can be realistically initiated for the deep Mediterranean, using the available technologies, protocols and monitoring programs and adapting the criteria, by neglecting the relevance of primary production, which could be replaced by the analysis of the inputs of primary organic matter and/or by starting from primary consumers and/or including the chemosynthetic primary production.

2.5. Descriptor 5: human-induced eutrophication

Eutrophication refers to “a process driven by enrichment of water by nutrients, especially compounds of nitrogen and/or phosphorus, leading to increased primary production and biomass of algae, changes in the balance of organisms, and water quality degradation” [131]. According to the MSFD, GES is achieved with respect to eutrophication when “Human-induced eutrophication is minimised, especially adverse effects thereof, such as losses in biodiversity, ecosystem degradation, harmful algae blooms and oxygen deficiency in bottom waters” (MSFD, 2008/56/EC, Annex I).

In coastal environments, eutrophication means the increase of primary production generally monitored through the chlorophyll-*a* (Chl *a*) content and/or macroalgal biomass [132–136]. The effects of eutrophication are the lowering of oxygen concentrations, losses of submerged aquatic vegetation, and mass mortalities especially at the sediment water interface [137,138]. The global occurrence and expansion of hypoxic/anoxic events at bathyal depths worldwide point to the need for better understanding and monitoring of the effects of such phenomena on deep-sea benthic communities [139,140]. Recent models foresee an oxygen decline from 1 to 7% in the next 100 years [141] with an increase in the extension of Oxygen Minimum Zones (OMZs) [142]. Farther, OMZs, with their naturally occurring low pH and oxygen, offer some hints as to the structure of deep-sea ecosystems affected by eutrophication [143,291].

Deep-sea ecosystems have been historically considered as a food-poor environment, and this is typically true for the deep Mediterranean Sea, especially in its eastern basin, but some areas may experience symptoms of eutrophication and oxygen depletion [144]. For instance, it has been reported that massive phytodetritus exports from highly productive coastal waters to the deep-sea floor [292]. Excessive C inputs in combination with the high bottom temperatures can cause episodic oxygen depletion in the deep sea [138,144]. Recent studies highlighted that deep-sea trophic status can be also affected by climate change, as the Western basin is expected to become more oligotrophic and the Eastern basin more eutrophic [145]. In addition, predicted increasing surface temperatures may affect water mass stratification and the formation of cold oxygenated deep water, modifying global ocean circulation and the dissolved oxygen availability in deep-water masses [146]. Local scale eutrophication could affect deep-sea sediments facing highly productive areas of the Mediterranean Sea, such as the Gulf of Lions, the northern Aegean Sea and the Ionian Sea receiving inputs from the Adriatic Sea.

However, the MSFD, in relation to the qualitative Descriptor 5, calls for an assessment of nutrients and organic matter inputs (Annex III of Directive 2017/845) and the use of the following criteria (Directive 2017/848): i) nutrient concentrations in the water column, ii) chlorophyll *a* in the water column, iii) harmful algal blooms, iv) photic limit, v) dissolved oxygen at the bottom of the water column, vi) opportunistic macroalgae, vii) macrophyte communities and viii) macrofaunal communities. These criteria can only be partially applied to the deep sea. Firstly, primary producers (e.g., macrophyte, macroalgae, harmful algal bloom) must be excluded, and the assessment of trophic status using variables measured in the water column can lead to misleading classifications [147]; see also [148]. Considering that oxygen depletion is one of the main causes of benthic faunal mortality, it is important to measure physical-chemical parameters and indicators also in the sediments [149]. In addition, the concentration of organic matter accumulated in surface sediments can provide a good indication of the eutrophication process occurring on the seafloor [147,150]. The current conceptual framework suggests the need to introduce new criteria and indicators, related to benthic ecosystems and, particularly, to the deep sea.

A group of core indicators is already utilised to monitor eutrophication in open waters, including: i) nutrients (nitrate, ammonium, phosphate), ii) dissolved oxygen and iii) phytoplankton (chlorophyll *a*, dominance). Zooplankton biomass is considered a potential, though not fully mature indicator [151] and references therein) because of

incomplete knowledge of its relationship to eutrophication. The monitoring of benthic ecosystems should include i) quantity and quality of sedimentary organic matter, and ii) biodiversity and taxonomic composition of benthic invertebrates [51,54,147,150,152]. Among indicators recently proposed to assess benthic trophic status of marine ecosystems, the quantity and biochemical composition of sedimentary organic matter has received the widest application, both in coastal and deep-sea ecosystems [150]; see also [148]. The concentrations of biopolymeric C (defined as the sum of C deriving from proteins, carbohydrates and lipids) and its algal fraction have been used to assess impacts of humans on benthic trophic status in different oceanic and coastal regions and varying water depths, within the Mediterranean basin [51, 147,150,152,153]. Changes in the quantity and quality of organic matter in the deep sea lead to responses from all benthic components, from prokaryotes to foraminifera, and from meiofauna to macrofauna [5]. Also the functional traits of macrofauna have been widely used as indicators of alteration and for measure the health status of marine benthic ecosystems [134]. Further, meiofauna could be considered a good indicator as it is highly sensitive to environmental changes, and particularly to organic enrichment due to eutrophication [152]. For these reasons, meiofauna have been recently proposed for the monitoring of eutrophication effects and for assessing the environmental quality of both coastal and deep-sea ecosystems [54,293].

2.6. Descriptor 6: sea floor integrity

Descriptor 6 requires that seafloor integrity is “at a level that ensures that the structure and functions of the ecosystems are safeguarded and benthic ecosystems, in particular, are not adversely affected” (Commission Decision 2010/477/EU). This involves addressing “physical damage, having regard to substrate characteristics”, and the “condition of benthic community”, the latter being directly related to Descriptor 1. The relevant pressures identified in the Commission Decision (EU) 2017/848 generically refer to “physical loss (due to permanent change of seabed substrate or morphology and to extraction of seabed substrate)” and to “physical disturbance to seabed (temporary or reversible)”. Four primary criteria address these points, three of which (D6C1 to D6C3) are specific for Descriptor 6, while two are also relevant for Descriptor 1 (D6C4 and D6C5).

The deep Mediterranean seafloor experiences two dominant physical disturbances associated with human activities: i) bottom-contact fisheries and ii) oil and gas activities [28,154,155,294]. Fisheries using bottom-contacting gear lead to direct alteration of seafloor morphology at large, medium and small scales [156,157]. Bottom trawling is a key driver for large-scale seascape change as it smoothens the natural topography [156]. Direct and indirect biological effects of bottom trawling have been demonstrated in terms of biogeochemical changes (e.g. less total amino acid concentration in sediments) and faunal desertification [51]. The Mediterranean Sea shows the highest fisheries footprint per unit landings in Europe [158], with peak intensities in the Tyrrhenian and the Adriatic Sea. In the Catalan margin, trawling impact is major down to 800 m depth [156]. Sediment resuspension from fishing grounds can propagate to wider and deeper areas eventually leading to suffocation and burial of VME [159]. Downslope moving gravity-driven resuspension flows enhance sedimentation rates far beyond fishing grounds, such as in canyon axes. Other types of fishing gear such as bottom touching longlines and gillnets could also have a significant adverse impact on vulnerable benthic communities and organisms such as black corals, gorgonians, scleractinians and many other habitat-forming species [105], because of breaking while pulling, ghost fishing or entanglement.

Activities undertaken by the offshore oil and gas industry may cause physical loss of the natural deep seabed. Physical (and chemical) impacts on the seafloor and subseafloor range from the installation of drilling rigs, wellheads and other structures on the seabed to the accumulation of litter including lost or abandoned equipment, consumables

and other materials. Today's deep-water (>200 m) oil and gas production in the Mediterranean Sea, or advanced prospects for it, takes place essentially offshore Egypt, Israel, Lebanon, Syria and Cyprus [160,161]. The environmental approach for the hydrocarbon industry in the Mediterranean Sea is developed in the Offshore Protocol of the Barcelona Convention, adopted in October 1994, which obliges countries to perform comprehensive EIAs after entering into force in December 2012. The EU adopted the Directive on Safety of Offshore Oil and Gas Prospection, Exploration and Production Activities in July 2013, which provides a blueprint of the best international practice also for non-EU countries in the Eastern Mediterranean that are new to the energy industry [162]. Further disturbance occurs in case of cable deployment, for not the cables and pipelines per se, rather for the impact of the anchoring of the supply vessel during the deployment of the cable.

Dumping of industrial waste in the deep Mediterranean Sea is a matter of concern for habitat integrity. Submarine canyons with heads close to the coast are favoured sites for direct deep-sea disposal [295]. Two aluminium-processing plants have discharged red mud waste in the deep Mediterranean Sea: one in France (Cassidaigne Canyon, Gulf of Lion) [163–165]; see also [148] and one in Greece (Gulf of Corinth, Antikyra Bay) [166–168]. Since 1988, Coal Fly-Ash (CFA) from the Hadera power plant, in Israel, has been dumped into a 16 km² disposal site some 70 km offshore, at a water depth of 1400 m, where a 0.5–1.0 cm thick ash layer has been noticed [169,170] together with severe impoverishment of benthic fauna. Israel allowed also long-term disposal of dredged sediments and industrial waste (1,900,000 m³) polluted with Hg, Cd, Pb, tributyltin and organotins, and PCBs at a site 1300 m deep [171].

Different proved tools is currently available to assess seafloor integrity. High-resolution maps of benthic substrata and habitats are increasingly required both to underpin environmental and socioeconomic impact assessments and to help in developing effective management measures [148,155,172–174]. Multibeam Echo-Sounders (MBES) and side scan sonars (SSS), map seabed areas with 100% spatial coverage at a resolution finer than 1 m², depending on the depth of data collection and on distance-to-bottom of the sensors [172]. Ground-truthing methods, such as the use of remotely operated vehicles (ROVs) and autonomous underwater vehicles (AUVs) [67,175], are widely available and could be applied according to the size and the nature of the area of interest [155,172,173,176]. Habitat suitability models try to predict the distribution of some habitats such as CWCs [68, 177–180]; [181]; [61,182,183]. However, because such models often include a large degree of uncertainty, decisions based entirely on modelling approaches may involve significant risk.

The revision of the MSFD (through the COMM. DEC. 2017/848/EU) emphasised that “Physical loss shall be understood as a permanent change to the seabed which has lasted or is expected to last for a period of two reporting cycles (12 years) or more”, but for this to be implemented, a very long time perspective is needed. All impacts described in this section have immediate effects (and sometimes also delayed effects) on seafloor communities, which in most cases could represent either a tipping point (e.g. large-scale seascape change) or require long time before any significant recovery could take place. A time-span of 12 years is possibly too short and it is urgent to proceed with a sound extensive evaluation of the current status of the deep benthic habitats in the Mediterranean Sea before human impact severely modifies or erases them from the face of our planet.

2.7. Descriptor 7: permanent alteration of hydrographical conditions

Descriptor 7 is geared towards addressing the problem of the permanent alteration of hydrographical conditions. These conditions are often affected by the presence of coastal infrastructure and other man-made activities (ports, artificial reefs, etc.). However, in most cases these structures impact coastal areas and only rarely can reach higher depths. Conversely, global climate change, combined with episodic

climate-driven events, can alter the “hydrographical conditions” also at depths. In recent decades, deep Mediterranean waters have experienced drastic changes resulting in an alteration of the stratification associated to temperature increases and salinity shifts [184,185]. At the end of the 1980s, climate change, changing hydrographic properties, surface circulation, and deep-water convection caused a ‘regime shift’ on a global scale [186] and throughout the Mediterranean basin [187]. During that period, the main site of deep-water formation shifted from the southern Adriatic to the Aegean sub-basins. This “Eastern Mediterranean Transient” (EMT; 1987–1994) event resulted in increased oxygen consumption [188,189] and, in the eastern Ionian, in a nutricline shoaling by about 150 m [190]. Other rapid hydrological changes have also occurred in the Western Mediterranean Sea. The “Western Mediterranean Transient” (WMT; 2004/05 and 2005/06) was characterised by the formation of warmer and denser new deep waters over the continental shelf as a result of cooling and evaporation of the surface layer and downslope cascading [19,184,191]. The high volumes of newly formed deep waters generated during intense cascading and convection events dramatically altered the hydrological structure of the basin, completely de-stratifying the water column and transferring massive heat and salt to the deep layers [19,184,192]. The 2004/05 event was the first of a series of similar events in the last decade that greatly altered the structure of the intermediate and, especially, the deep layers of the Western Basin [22]. Cascading events transport huge amounts of nutrients and organic matter, to bathyal depths [5,19,193,194]. Hydrographic preconditioning (heat and salt content and structure of the water column before the onset of convection), and atmospheric forcing (heat, freshwater and buoyancy fluxes) triggered deep-water formation [148]. Moreover, progressive increase in heat and salt content in the intermediate layer, advected from east to west, favoured new dense water formation in the North-Western Mediterranean basin. Multiple heat and saline anomalies characterised the Mediterranean Sea from 1950 to 2000 [195,196] and although these alterations cannot be considered permanent, all of these changes have long-term effects.

The multidisciplinary Mediterranean Targeted Projects MTP-I and MTP-II/MATER (1993–2000; [197], the MEDAR/MEDATLAS database [198], the SeaDataNet (i.e., <https://www.seadatanet.org/>) and EMODnet (<http://www.emodnet-physics.eu>) infrastructures created datasets on temperature, salinity oxygen, silicate, nitrates and phosphates, but in some cases with insufficient coverage in the eastern region (including Tunisia, Libya, Croatia and Turkey; [199].

Since the 2000's, national and international programmes (e.g. EU-PERSEUS and MedSeA, IT-VECTOR, FR-MERMEX, E-RADMED) produced hydrological data on the whole Mediterranean Basin. Regular cruises, mooring lines, and deployment of new instruments and infrastructure (Argo floats, gliders) now support intensive collection of in situ observations in the North-Western Mediterranean Sea. Argo floats, autonomous profiling floats, drift at a given depth for a given time period. After drifting for a set time, they sink to 2000 m and profile temperature and salinity during the upcast. These data proven useful in describing deep-water formation [200]. Moreover, in the last decade, glider technology mainly in the North-Western Mediterranean Sea has enabled repeated cross-basin transects at depths from the surface to 1000 m. In addition, numerical models implemented at regional or local scales use these data to elucidate water mass formation and spreading, and basin-scale hydrological dynamics [201,202].

Long-term monitoring of basic hydrological parameters (temperature and salinity), collected as time series with appropriate temporal resolution (i.e. sampling intervals that resolve all relevant timescales) represent a science priority in the context of climate change study for key locations in the Mediterranean Sea (e.g. straits and channels, zones of dense water formation and spreading, deep basins) [32,203]. The HYDROCHANGES network aims to address this need by deploying moorings fitted with Conductivity, Temperature, Depth sensors (CTDs) at key locations for monitoring temperature and salinity [203]. The FixO3 (Fixed point Open Ocean Observatory network, <http://earthvo>.

fixo3.eu/) programmes and the EMSO (European Multidisciplinary Seafloor and water column Observatories, <http://emso.eu/>) EU Infrastructure provide additional near-bottom data from fixed-point observations. For example, DYFAMED and LION deep-water stations, and ANTARES neutrino telescope in the North-Western Mediterranean or the KM3NET in the Ionian Sea [204,205], continuously monitor sets of specific parameters. Long time series provided by these mooring stations have contributed pivotal findings on the deep dynamics of the Mediterranean Sea in last years. In order to partially interpolate data within homogeneous habitats, scientists have generated gridded products through objective analysis of available observations (such as numerical models with data assimilation delivered by Copernicus downstream services (<http://marine.copernicus.eu/services-portfolio/access-to-products/>)).

2.8. Descriptor 8: concentrations of contaminants giving rise to pollution effects

The input of xenobiotic substances represent one of the major threats for ocean health [206]. Hydrophobic pollutants, such as organo-halogenates and polycyclic aromatic hydrocarbons (PAH), enter the marine environment through effluent discharges, atmospheric deposition, runoff and other means [207]. Once in the water column, the adsorption onto particulate matter transfers these compounds from the surface to the deep waters and sediments [296]. Particle settling is also favoured by biological processes, and by lateral transport from continental shelves [208–210]. DSWC is a massive mechanism of pollutant transfer to the open deep ocean [19]. Higher fluxes of organohalogen pollutants and Polycyclic Aromatic Hydrocarbons (PAH) occur during these cascading events [211]. PAH settling fluxes in the north-western Mediterranean Sea vary widely [212,213], with highest concentrations of contaminants in the Alboran Sea [214], and much lower values in Sardinia and in the Southern Ionian Sea [215,216]. However, these values greatly exceed atmospheric deposition of PAH in central sites of the Western Mediterranean Sea, thus highlighting the role of river discharge [208,217,218]. Qualitative differences are also observed in relation to these transfer processes. Sediments of coastal areas, continental shelves and slopes have higher proportions of petrogenic PAHs whereas the deep basin of the north-western Mediterranean Sea is characterised by high amounts of pyrogenic PAHs.

Organochlorine compounds such as Polychlorobiphenils (PCBs) and chlorinated pesticides, characterised a group of Persistent Organic Pollutants (POPs) of worldwide concern due to their toxic effects [219]. Notwithstanding the discontinued use of these compounds in most world areas, thanks to relevant national regulations and international agreements such as the Stockholm Convention, their extensive occurrence is still observed. Their high lipophilicity, hydrophobicity, chemical stability and resistance to biological degradation have led to their accumulation in biological tissues and biomagnification through the food chain.

Radioactive compounds in the Mediterranean Sea are derived from the fallout of nuclear weapon testing and the Chernobyl accident. In sediments, concentrations of ^{137}Cs and $^{239+240}\text{Pu}$ have been measured in various parts of the Mediterranean Sea, including deep basins. The concentrations are generally higher in coastal ecosystems because land-based sources can exceed atmospheric inputs [210,220]. Concentrations in biota are presently undistinguishable from those in areas without point sources. Hence, the relevance of these contaminants lies in their usefulness as process tracers, more than on their impact on the environment. Studies on radionuclides in marine organisms also underscore that the radionuclide levels are constantly decreasing due to modifications of the inputs. Very little work has been done to examine the trophic transfers of man-made radionuclides [221]. Finally, neglected impacts that can be very important in several areas of world are military activities. Information on their impacts on the environment are relatively scarce and are often studied after several years from their production

and without any baseline available [33,222].

Atmospheric inputs constitute one of the major sources of Trace Elements (TE) to the deep Mediterranean Sea [73,223], where TE concentrations in waters are typically higher than in other areas of the world ocean. In addition, Cd, Cu and Ni (as well as Cr) are dominated by lateral advection and vertical mixing rather than by biogeochemical cycling [224]. The hydrologic regime of the Mediterranean Sea tends to transfer the pollutants and nutrients to the Atlantic by bottom water flow transport. Our knowledge on the concentrations, fluxes, and behaviour of trace elements, radionuclides and organic substances in the deep waters and sediment and their toxicological impacts on habitats and organisms is scarce [220]. Pollutants with hydrophobic properties, e.g. PCBs and mercury, accumulate in biota and thus in the food web. In case of chronic pollution events, the concentration of contaminants should be analysed in sediments collected by sediment cores, which enable the reconstruction of temporal trends. Sample collection of water at different depths and analysis of the dissolved and particulate matter would also be important. Abundance of populations and estimates of the extent of habitats adversely affected by chronic pollution should be assessed concurrently.

Nonetheless, information on pollutants in the deep sea is almost completely lacking, and this represent the main gap in the application of the criteria needed to determine the D8.

2.9. Descriptor 9: contaminants in fish and other seafood for human consumption

Descriptor 9 focuses on the accumulation of toxic, persistent and liable substances in wild deep-sea organisms used for human consumption (i.e., mostly teleost and decapod crustaceans) and the contaminants considered by D9 are only part of those of interest for the D8 (cfr. Regulation EC 1881/2006 and its amendments [225,226]). Each Member State may ignore specific contaminants and/or include additional ones [227,228]. In any case the monitoring of the contaminants accumulated in the deep-sea biota should at least consider the following compounds for which regulatory levels have been set: i) heavy metals (lead, cadmium and mercury); ii) PAHs; iii) dioxins (including dioxin-like PCBs). In addition, the following contaminants of relevance should be monitored: i) non-dioxins like PCBs; ii) phthalates; iii) organochlorine pesticides; iv) organotin compounds; v) brominated flame retardants; vi) polyfluorinated compounds. Also, artificial radionuclides should be monitored in case of nuclear accidents or any other radioactive emergencies that could lead to or has led to significant radioactive contamination of food.

Contaminants in fish and other seafood might derive from numerous anthropogenic sources described for the D8. Chemical contamination in fish and seafood results from a complex process that balances inputs of contaminants, mostly through diet, and their excretion [124,229,230].

Investigating contamination levels in fish and seafood requires understanding which contaminants exceed regulatory limits, how much they alter food webs, and what metabolic processes are involved in detoxification. The presence of xenobiotics in the deep Mediterranean organisms has been repeatedly documented [231,232] with deep-sea Mediterranean fishes tending to exhibit higher levels of metal accumulation than those of populations inhabiting other areas such as the Atlantic Ocean [233]. Red-shrimps, *Aristeus antennatus* and *Aristaeomorpha foliacea* may be useful indicator species of levels of deep-sea contamination (e.g., see data on *A. antennatus*, [234]). Contaminants in fish muscle and liver have been investigated in the most abundant deep-sea megafaunal species, e.g. *Alepocephalus rostratus*, *Coelorhynchus mediterraneus*, *Coelorhynchus caelorhynchus*, *Trachyrhynchus trachyrhynchus* and *Nezumia sclerorhynchus*, *Chimaera monstrosa*, *Lophius budegassa*, *Lepidion lepidion* [235], revealing mercury concentration exceeding $0.5 \mu\text{g g}^{-1}$ muscle wet weight in all species but one. This represents the threshold value indicated by the European Commission as acceptable for human consumption. High mercury concentration is a distinct feature of the

Mediterranean Sea (the so-called “Mediterranean mercury anomaly”; [236,237]. The Hg concentration is even higher in some deep-sea fish species [124,235,238]. Most deep-sea species are long-lived and slow-growing, which favours the bioaccumulation of pollutants [235, 239–241]. Since some deep-sea species are of commercial interest, the high contamination level poses serious risks for human health [242, 243]. Despite this, few studies have investigated the concentrations of contaminants in the deep Mediterranean fauna fished for human consumption [124,235,244,245]. In this regard, the Gulf of Lion is the best-investigated area, whereas the Levantine Basin (with the exception of Israel waters [231]; are the least studied. A comparison of the concentrations of xenobiotics in fish collected in the NW Mediterranean Sea, in 1996 and 2009 [229,240,246] indicate that their contamination did not change over time.

The application of the criteria of Descriptor 9 should consider the concentration, the thresholds, and the contamination sources. Regulators should also consider the species of interest for human diets and their ability to bioaccumulated pollutants. GES would be achieved if all contaminants occur at levels below those established for safe human consumption.

2.10. Descriptor 10: marine litter

Two primary and two secondary criteria are associated to Descriptor 10: i) the composition, amount and spatial distribution of litter (D10C1) and of micro-litter (D10C2) “on the coastline, in the surface layer of the water column, and on the seabed, are at levels that do not cause harm to the coastal and marine environment” (primary) and ii) the amount of litter ingested by marine animals, which should not reach a level that adversely affect the health of the species (D10C3) and the number of individuals which are adversely affected due to litter, such as by entanglement, other types of injury or mortality, or health effects (D10C4) (secondary). Each sub-region should assess the outcomes for all criteria and as well as threshold values.

Marine litter represents a threat for the health of the deep Mediterranean Sea due to its limited exchange with other basins, dense population, touristic and industrialized coastlines, and heavy maritime traffic [297]. The sources of marine litter to the deep-sea floor of the Mediterranean Sea are either from land (river discharge, storm drains, sewage treatment plants and industrialized areas) or marine (fishing activities, commercial and recreational shipping, aquaculture, direct dumping), and include plastics (accounting for >70% of the total), glass, metal, clinker, cardboard and fabrics [27,29,67,247–249,297]. The quantity and composition of marine litter differs among regions and changes with depth, probably as a result of a complex set of interactions between hydrodynamics, geomorphology, and anthropogenic sources [27,249,297]. The abundance of marine litter items in the deep Mediterranean Sea varies from 500 items km^{-2} on the continental slopes off Malta and Cyprus (Mifsud et al., 2013; [250], the Tyrrhenian Sea [251], or the Adriatic Sea [247], to more than 2000 items km^{-2} in the Antalya Bay in the Eastern Mediterranean or in the submarine canyons of the Gulf of Lion and of the Catalan Sea [27,247]. Astonishingly high litter abundance of up to 1.3 million of items km^{-2} were reported at 300–600 m depth in the Messina Strait canyons (Central Mediterranean Sea) [252]. Litter abundance found in submarine canyons and depths greater than 500 m typically exceeds that at shallower depths, suggesting that submarine canyons can act as primary conduits of litter from the coast to the deep sea [27,247]. Superposition of highly efficient source-to-sink sedimentary transport (with flash-flood generated hyperpycnal flows) and strong urbanization of the coastal area promote the occurrence of large litter hotspots in the deep sea [252]. In addition to large marine debris, concern has grown about microplastics (i.e., <1–5 mm in diameter; [253], which can directly enter the ocean also through cosmetic abrasives (i.e. microbeads), preproduction plastic pellets, or textile fibres known as primary plastics. Additionally, combined mechanical, biological, photic and thermal actions can break

down larger plastic objects into numerous small fragments, which are defined as secondary microplastics. Depending on the density of the polymer, microplastics may sink and behave as very fine-grained sediments (for example polyester; [254], or they may float and subsequently sink following colonization by organisms, adsorption to phytoplankton, and/or aggregation with organic debris. Fibres appear to dominate the microplastics reported in Mediterranean deep-sea sediments [254–256]. Recent studies indicate that the Mediterranean deep-sea floor might act as a long-term sink for microplastics where the abundances of such particles can exceed those in surface waters [256]. Marine litter in the Mediterranean deep sea may significantly affect different ecological compartments and, consequently, human health, with potentially severe economic impacts. Biotic effects of large and small items include entanglement, ingestion, colonization and rafting [26,27,248,249,251, 257–259]. However, information on the actual effects of (micro)plastics on deep-sea organisms and trophic webs is still limited [260].

Marine litter in deep sea produces economic impacts primarily on the fishery sector, damaging vessels and fishing equipment due to entanglement of catch, loss of target species through ghost fishing, or reduced reproductive capacity of benthic organisms consuming microplastics [261]. Furthermore, marine litter may contain pollutants (hazardous plastic additives, POPs) that exert toxic and endocrine disruptive effects on marine organisms that ingest plastics [262].

2.11. Descriptor 11: introduction of energy including underwater noise

Descriptor 11 deals with introduction of energy into the marine environment. Underwater noise can be pulsed or continuous. MSFD currently focuses on two criteria: anthropogenic pulsed (D11C1) and continuous low-frequency (D11C2) sounds in water. D11C1 addresses the space-time distribution of pulsed noise sources, whereas D11C2 addresses levels of continuous noise, using in situ measurements and models. Pulsed noise may cause direct acute effects such as hearing loss, tissue damage, and death of individuals of sensitive species such as cetaceans. Whereas continuous or chronic noise exposure mainly causes stress and behavioural alterations, with negative effects on deep-sea organisms [263]. The proposed strategy on noise monitoring recommends several adaptations in the case of the deep Mediterranean. Particularly, both indicators are closely related to the acoustic biology of deep-diving marine mammal species, such as sperm whale and Cuvier’s beaked whale. Pulsed noise can be monitored by setting up a register of anthropogenic activities, reporting on date, location, proportion of days within a given period and over a given geographical scale in which activities generating pulsed sounds occur. This could be done through the deployment of hydrophones (e.g., permanent or semi-permanent PAM) on new infrastructures (or implementation of the existing infrastructures) and their subsequent future development into larger geographic network: [264].

A variety of phenomena generates noise in the ocean, either from physical processes, such as wind-generated waves, earthquakes, precipitation, or from biological phenomena such as whale songs, dolphin clicks, and fish vocalizations [265]; not all reach the deep sea. Fish produce sounds for their navigation, habitat selection and mating, as well as to communicate [266]. Marine mammals use sound as a primary tool for underwater communication [267], mating and social interaction [268], and for tracking the prey [269].

Anthropogenic noise can reach the deep sea, through commercial shipping, oil and gas exploration, fishing, and scientific research; all of these sources currently contribute to the general noise budget of the ocean [265]. The impact of noise on marine is being increasingly investigated [270,271]. Noise sources are divided into three frequency bands: low (10–500 Hz), medium (500 Hz–25 kHz) and high (>25 kHz). Anthropogenic sources dominate the low-frequency band, and include commercial shipping and seismic emissions for hydrocarbon exploration. Minimal attenuation of low-frequency sound allows long-distance propagation. Sea-surface agitation (breaking waves, spray, bubble

formation and collapse, and rainfall) and various sonars (e.g. military and multibeam seabed mapping), as well as small vessels produce most medium frequency sound. Greater attenuation limits propagation of noise in the mid-frequency band over long distances, and only local or regional (10s of km distant) sound sources contribute to this ambient noise field. At high frequencies, extreme acoustic attenuation confines all noise sources to the area close to the receiver.

Oil industry operations have traditionally focused on shallow, continental shelf waters, but exploration is moving in deeper waters (>500–1000 m). Expansion of oil exploration into deeper water has increased the potential for long-range propagation of seismic reflection signals. Indeed, sound in deep waters can propagate greater distances than in shallow-water ecosystems, by moving through the deep sound channel [271]. Seismic surveys currently target all regional seas in the south-eastern Mediterranean, apart from the Aegean Sea [272]. This expansion stresses the transboundary aspect of seismic surveys and calls for international cooperation.

Anthropogenic noise can cause physical and biological damage, such as behavioural changes and stress, especially in marine mammals, sea turtles and fish [273,274]. The occurrence of low frequency noise in the deeper part of the basins is particularly important for deep-diving marine mammals, such as toothed whales, because the ambient noise they use as a background for echolocation decreases rapidly with depth [275–277]. Indeed, small odontocetes produce high frequency sounds (ranging from 70 kHz to more than 150 kHz), while sperm whales, *Physeter macrocephalus*, during diving, make sound with frequencies ranging to more than 30 kHz which are detectable within 10–15 km. The fin whale *Balaenoptera physalus*, the only mysticete constantly present in the Mediterranean Sea, emits mostly infrasonic signals (20–40 Hz), which are emitted in long sequences and can be detected at large distances.

Research on sea turtles in the South-Eastern Mediterranean region revealed that they can detect low frequency sounds that overlap with seismic airgun frequencies, these are high-intensity, low-frequency impulsive noise at regular intervals, mostly between 10 and 300 Hz [278,279]. Airguns can stress the sea turtles, *Caretta caretta* [280]. The impacts of anthropogenic noise on sharks and rays are poorly studied, with most research to date focusing outside the Mediterranean region [281]. Fish sensitivity to certain frequencies varies among species [278]. Recent studies demonstrate negative effects of seismic survey airgun operations even in zooplankton [282].

In situ acoustic measurements can document continuous low-frequency sound, gathering field data on ambient noise in a given location. Understanding the large-scale influence of artificial noise on marine organisms and ecosystems represents the main gap to the application of the D11 on the deep sea. Deep-sea observatories offer new opportunities to assess the presence and effects of noise in on deep-sea life [32]. Deep-sea cabled observatories (i.e. NEMO-SN1 in the Western Ionian Sea, [283,284]; ANTARES in the Ligurian Sea, [285]; and PYLOS in the South Ionian Sea, <http://www.fixo3.eu/observatory/pylos/>) are equipped with hydrophones for passive acoustic monitoring. Besides these measurements, and especially for monitoring continuous low-frequency sound in deep sea, modelling approaches (both for single sources or distributed sources of noise, from the most advanced Dynamic Ambient Noise Prediction System elaborated by the U.S. for modelling multiple sources, to the Acoustic Integration Model used for modelling the effects of noise on cetaceans [286]; may reduce the time required to establish trends and patterns.

3. Future implementation

In order to effectively implement the deep-sea MSFD, we need to identify the criteria to achieve or maintain GES in open waters and deep-sea bottoms, including “*spatial protection measures, contributing to coherent and representative networks of marine protected areas, adequately covering the diversity of the constituent ecosystems, such as special areas of*

conservation pursuant to the Habitats Directive, special protection areas pursuant to the Birds Directive, and marine protected areas as agreed by the Community or Member States concerned in the framework of international or regional agreements to which they are parties” (MSFD, 2008/56/EC, Article 13).

The MSFD takes an overarching and integrated approach by focusing on achieving GES and targets, and we therefore recommend exploring and assessing synergies between the different treaties, directives, and conventions (e.g. see Descriptor 6 section) so that, wherever possible, the programme of measures and proposed MSFD monitoring simultaneously address the requirements of other legislations.

Our analysis indicates that the 11 Descriptors promulgated by the MSFD (MSFD, 2008/56/EC) can be adapted and applied to the deep sea. Several Descriptors (D1, D2, D3, D6, D8, D10) can be readily implemented, others (D4, D9 and D11) require additional data in order to set up benchmark and threshold values, while two (D5, D7) require changes in the assumptions and/or modification in the concept of “permanent”.

Priority ecological variables, spatial distribution, extent of pressures and impacts ought to be identified and standardized in order to establish targets and indicators addressing the distinct conditions in the deep sea. The expertise, tools and resources required for deep-sea monitoring are not universally available to all Mediterranean Member States (MS), nor to countries in the southern and easternmost Mediterranean. These limitations may be overcome by initiating deep sea MSFD-focused monitoring in already data-rich locations (presumably off MS), and pioneering joint-effort monitoring in collaboration with non-MS, to enhance awareness, capacity building, and gain much needed data on scantily studied regions. Given the costs entailed by scientific and technical expertise, tools and infrastructure required for deep-sea research and monitoring, we advocate for EU-level financial support for MS/non-MS collaboration spanning joint fieldwork, training (early career research fellowships, workshops) and public awareness communication.

Since the millennium, increased awareness of the vulnerability of deep-sea ecosystems has changed attitudes concerning their protection and conservation [146]. Yet, for effective regulatory measures, legislators and managers require scientific evidence, which follows from basic scientific research and monitoring. Ecosystem-based management of the Mediterranean deep sea pressingly requires comprehensive analysis of available data, new data from yet unexplored regions, and impact assessment studies. Mounting evidence points to the vulnerability of the deep biota to anthropogenic disturbance that may result in biodiversity loss, urgent implementation of the MSFD in the Mediterranean deep sea will go a long way towards conserving its unique biodiversity and habitats.

Acknowledgements

This study has been supported by the DG ENV project IDEM (Implementation of the MSFD to the Deep Mediterranean Sea; contract EU No 11.0661/2017/750680/SUB/EN V.C2). MC and AS-V from University of Barcelona acknowledge support from the Spanish government through Red BAMAR (ref.: CGL2016-81854-REDT), a network on marine litter, and RTD projects NUREIEV (ref. CTM2013-44598-R) and NUREIEVA (ref. CTM2016-75953-C2-1-R) on far-field and near-field impacts of the Portman Bay, SE Spain, coastal submarine mine tailings disposal site. Generalitat de Catalunya autonomous government funding to CRG Marine Geosciences (ref. 2017 SGR 315) within its support scheme to excellence research groups is equally acknowledged.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.marpol.2019.103781>.

References

- [1] A.R. Longhurst, Chapter 1 - toward an ecological geography of the sea, in: A. R. Longhurst (Ed.), *Ecological Geography of the Sea*, second ed., Academic Press, 2017, pp. 1–17.
- [2] M. Taviani, The Mediterranean benthos from late Miocene up to present: ten million years of dramatic climatic and geological vicissitudes, *Biol. Mar. Mediterr.* 9 (2002) 445–463.
- [3] R. Danovaro, C. Corinaldesi, G. D'Onghia, B. Galil, C. Gambi, A.J. Gooday, N. Lampadariou, G.M. Luna, C. Morigi, K. Olu, P. Polymenakou, Deep-sea biodiversity in the Mediterranean Sea: the known, the unknown, and the unknowable, *PLoS One* 5 (8) (2010) 11832.
- [4] C.C. Emig, P. Geistdoerfer, The Mediterranean deep-sea fauna: historical evolution, bathymetric variations and geographical changes, *Carnets de Geologie, Madrid* 4 (A01) (2004) 10, <https://doi.org/10.4267/2042/3230>.
- [5] R. Danovaro, A. Dinet, G. Duineveld, A. Tselepidis, Benthic response to particulate fluxes in different trophic environments: a comparison between the Gulf of Lions–Catalan sea (western mediterranean) and the cretan sea (Eastern-Mediterranean), *Prog. Oceanogr.* 44 (1999) 287–312.
- [6] N. Bianchi, C. Morri, Marine biodiversity of the Mediterranean Sea: situation, problems and prospects for future research, *Mar. Pollut. Bull.* 40 (2000) 367–376.
- [7] R. Danovaro, A. Pusceddu, Ecomanagement of biodiversity and ecosystem functioning in the Mediterranean Sea: concerns and strategies, *Chem. Ecol.* 23 (2007) 347–360.
- [8] M. Coll, C. Piroddi, K. Kaschner, F. Ben Rais Lasram, J. Steenbeek, et al., The biodiversity of the Mediterranean Sea: status, patterns and threats, *PLoS One* 5 (8) (2010), e11842, <https://doi.org/10.1371/journal.pone.0011842>.
- [9] C. Lejeune, P. Chevaldonné, C. Pergent-Martini, C. Boudouresque, T. Pérez, Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea, *Trends Ecol. Evol.* 1204 (2010), <https://doi.org/10.1016/j.tree.2009.10.1009> published online.
- [10] E. Ramirez-Llodra, J.B. Company, F. Sardá, G. Rotllant, Megabenthic diversity patterns and community structure of the Blanes submarine canyon and adjacent slope in the Northwestern Mediterranean: a human overprint? *Mar. Ecol.* 31 (2010) 167–182.
- [11] K. Olu-Le Roy, M. Sibuet, A. Fiala-Médioni, S. Gofas, C. Salas, A. Mariotti, J. P. Foucher, J. Woodside, Cold seep communities in the deep eastern Mediterranean Sea: composition, symbiosis and spatial distribution on mud volcanoes, *Deep Sea Res. Part I* 51 (2004) 1915–1936.
- [12] M. Taviani, The deep-sea chemoautotroph microbial world as experienced by the Mediterranean metazoans through time, *Lect. Notes Earth Sci.* 131 (2011) 277–295.
- [13] M. Taviani, Marine chemosynthesis in the Mediterranean sea, in: S. Goffredo, Z. Dubinsky (Eds.), *The Mediterranean Sea: its History and Present Challenges*, Springer Science+Business Media, Dordrecht, 2014, pp. 69–83, https://doi.org/10.1007/978-94-007-6704-1_5, 2014.
- [14] U. Fernandez-Arcaya, E. Ramirez-Llodra, J. Aguzzi, A.L. Allcock, J.S. Davies, A. Dissanayake, P. Harris, K. Howell, V.A.I. Huvenne, K. Ismail, M. Macmillan-Lawler, J. Martín, L. Menot, M. Nizinski, P. Puig, A. Rowden, F. Sanchez, H. A. Steward, I. Van den Beld, Ecological role of submarine canyons and need for canyon conservation: a review, *Front. Mar. Sci.* 31 (2017).
- [15] EEA, State and Pressures of the Marine and Coastal Mediterranean Environment, Environmental Issue series, No5, European Commission, 1999, p. 137.
- [16] R. Danovaro, Pollution threats in the Mediterranean sea: an overview, *Chem. Ecol.* 19 (1) (2003) 15–32.
- [17] WWF/IUCN, The Mediterranean Deep-Sea Ecosystems: an Overview of Their Diversity, Structure, Functioning and Anthropogenic Impacts, with a Proposal for Conservation, IUCN, Malaga and WWF, Rome, 2004.
- [18] UNEP, Implementation of the Ecosystem Approach (EcAp) in the Mediterranean by the Contracting Parties in the Context of the Barcelona Convention for the Protection of the Marine Environment and the Coastal Region of the Mediterranean and its Protocols, 2012. http://www.rac-spa.org/ecapmed_i.
- [19] M. Canals, P. Puig, X. Durrieu de Madron, S. Heussner, A. Palanques, J. Fabres, Flushing submarine canyons, *Nature* 444 (2006) 354–357.
- [20] C. Ulses, C. Estournel, P. Puig, X. Durrieu de Madron, P. Marsaleix, Dense water cascading in the northwestern Mediterranean during the cold winter 2005. Quantification of the export through the Gulf of Lion and the Catalan margin, *Geophys. Res. Lett.* 35 (2008) L07610, <https://doi.org/10.1029/2008GL03325>.
- [21] A. Sanchez-Vidal, M. Canals, A.M. Calafat, G. Lastras, R. Pedrosa-Pàmies, M. Menéndez, R. Medina, J.B. Company, B. Hereu, J. Romero, T. Alcoverro, Impacts on the deep-sea ecosystem by a severe coastal storm, *PLoS One* 7 (2012), e30395, <https://doi.org/10.1371/journal.pone.0030395>.
- [22] X. Durrieu de Madron, L. Houpert, P. Puig, A. Sanchez-Vidal, P. Testor, A. Bosse, C. Estournel, S. Somot, F. Bourrin, M.N. Bouin, M. Beauverger, L. Beguery, A. Calafat, M. Canals, C. Cassou, L. Coppola, D. Dausse, F. D'Ortenzio, J. Font, S. Heussner, S. Kunesch, D. Lefevre, H. Le Goff, J. Martín, L. Mortier, A. Palanques, P. Raimbault, Interaction of dense shelf water cascading and open-sea convection in the northwestern Mediterranean during winter 2012, *Geophys. Res. Lett.*, *Am. Geophys. Union* 40 (2013) 1379–1385, 2013.
- [23] M. Taviani, L. Angeletti, L. Beuck, E. Campiani, S. Canese, F. Fogliani, A. Freiwald, P. Montagna, F. Trincardi, Reprint of 'On and off the beaten track: megafaunal sessile life and Adriatic cascading processes, *Mar. Geol.* 375 (2016) 146–160.
- [24] A.D. Rogers, The biology of *Lophelia pertusa* (Linnaeus 1758) and other deep-water reef-forming corals and impacts from human activities, *Int. Rev. Hydrobiol.* 84 (4) (1999) 315–410.
- [25] J.A. Koslow, G.W. Boehlert, J.D.M. Gordon, R.L. Haedrich, P. Lorange, N. Parin, Continental slope and deep-sea fisheries: implications for a fragile ecosystem, *ICES (Int. Council. Explor. Sea) J. Mar. Sci.* 57 (2000) 548–557.
- [26] M. Bo, S. Bava, S. Canese, M. Angiolillo, R. Cattaneo-Vietti, G. Bavestrello, Fishing impact on deep Mediterranean rocky habitats as revealed by ROV investigation, *Biol. Conserv.* 171 (2014) 167–176.
- [27] X. Tubau, M. Canals, G. Lastras, X. Rayo, J. Rivera, D. Amblas, Marine litter on the floor of deep submarine canyons of the Northwestern Mediterranean Sea: the role of hydrodynamic processes, *Prog. Oceanogr.* 134 (2015) 379–403.
- [28] G. D'Onghia, C. Calulli, F. Capezzuto, R. Karlucci, A. Carluccio, A. Grehan, A. Indennitate, P. Maiorano, F. Mastrototaro, A. Pollice, T. Russo, Anthropogenic impact in the Santa Maria di Leuca cold-water coral province (Mediterranean Sea): observations and conservation straits, *Deep Sea Res. Part II Top. Stud. Oceanogr.* 145 (2017) 87–101.
- [29] A. Mecho, J. Aguzzi, B. De Mol, G. Lastras, E. Ramirez-Llodra, N. Bahamon, J. B. Company, M. Canals, Visual faunistic exploration of geomorphological human-impacted deep-sea areas of the north-western Mediterranean Sea, *J. Mar. Biol. Ass. UK* 98 (2017) 1241–1252.
- [30] P.G. Ryan, C.J. Moore, J.A. van Franeker, C.L. Moloney, Monitoring the abundance of plastic debris in the marine environment, *Philos. Trans. R. Soc. Biol. Sci.* 364 (2009) 1999–2012.
- [31] R. Danovaro, L. Carugati, A. Boldrin, A. Calafat, M. Canals, J. Fabres, K. Finlay, S. Heussner, S. Miserocchi, A. Sanchez-Vidal, Deep-water zooplankton in the Mediterranean Sea: results from a continuous, synchronous sampling over different regions using sediment traps, *Deep Sea Res. Part I* (2017) 103–114.
- [32] J. Aguzzi, D. Chatzievangelou, S. Marini, E. Fanelli, R. Danovaro, S. Flögel, N. Lebris, F. Juanes, F. De Leo, J. Del Rio, L.S. Thomsen, C. Costa, G. Riccobene, C. Tamburini, D. Lefevre, C. Gojak, P.M. Poulain, P. Favali, A. Griffa, A. Purser, D. Cline, D. Edgington, J. Navarro, S. Stefanni, J.B. Company, New high-tech interactive and flexible networks for the future monitoring of deep-sea ecosystems, *Environ. Sci. Technol.* 53 (12) (2019) 6616–6631.
- [33] R. Danovaro, E. Fanelli, J. Aguzzi, D. Billett, L. Carugati, C. Corinaldesi, A. Dell'Anno, C. Gjerde, A.J. Jamieson, S. Kark, C. McClain, L. Levin, N. Levin, E. Ramirez-Llodra, H. Ruhl, C.R. Smith, P.V.R. Snelgrove, L. Thomsen, C. Van Dover, M. Yasuhara, Ecological variables for developing a global deep-ocean monitoring and conservation strategy, *Nat. Ecol. Evol.* (2019) in press.
- [34] S. Raicevich, P. Battaglia, T. Fortibuoni, T. Romeo, O. Giovanardi, F. Andaloro, Critical inconsistencies in early implementations of the marine strategy framework directive and Common fisheries policy objectives hamper policy synergies in fostering the sustainable exploitation of mediterranean fisheries resources, *Front. Mar. Sci.* 4 (2017) 316.
- [35] P.V.R. Snelgrove, K. Soetaert, M. Solan, S. Thrush, C.-L. Wei, R. Danovaro, R. W. Fulweiler, H. Kitazato, B. Ingole, A. Norikko, R.J. Parkes, N. Volkenborn, Contrasting biogeochemical and biological estimates of carbon turnover on the global seafloor, *Trends Ecol. Evol.* 33 (2018) 96–105.
- [36] J.L.S. de Vivo, J.C. Rodríguez Mateos, Marine governance in the Mediterranean sea, in: M. Gilek, K. Kern (Eds.), *Governing Europe's Marine Environment: Europeanization of Regional Sea or Regionalization of EU Policies?* 2015, Ashgate Publishing, 2015. Published by.
- [37] E. Fanelli, J.E. Cartes, V. Papiol, Assemblage structure and trophic ecology of deep-sea demersal cephalopods in the Balearic basin (NW Mediterranean), *Mar. Freshw. Res.* 63 (2012) 264–274.
- [38] E. Fanelli, J.E. Cartes, V. Papiol, C. López-Pérez, Environmental drivers of megafaunal assemblage composition and biomass distribution over mainland and insular slopes of the Balearic Basin (Western Mediterranean), *Deep Sea Res., Part I* 78 (2013) 79–94.
- [39] A. Quetglas, F. Ordines, M. Gonzale, N. Zaragoza, S. Mallol, M. Valls, A. De Mesa, Uncommon pelagic and deep-sea cephalopods in the Mediterranean: new data and literature review, *Mediterr. Mar. Sci.* 14 (2013) 69–85.
- [40] J. Bertrand, L. De Sola, C. Papaconstantinou, G. Relini, A. Souplet, The general specifications of the MEDITS surveys, *Sci. Mar.* 66 (2002) 9–17.
- [41] J. Aguzzi, N. Bahamon, Modeled day-night biases in decapod assessment by bottom trawling survey, *Fish. Res.* 100 (2009) 274–280.
- [42] J. Aguzzi, J.B. Company, N. Bahamon, M.M. Flexas, S. Tecchio, U. Fernandez-Arcaya, J.A. García, A. Mechó, S. Koenig, M. Canals, Seasonal bathymetric migrations of deep-sea fishes and decapod crustaceans in the NW Mediterranean Sea, *Prog. Oceanogr.* 118 (2013) 210–221.
- [43] P. Vasilakopoulos, C.D. Maravelias, G. Tserpes, The alarming decline of Mediterranean fish stocks, *Curr. Biol.* 24 (2014) 1643–1648.
- [44] M. Cardinale, G.C. Osio, G. Scarcella, Mediterranean Sea: a failure of the European fisheries management system, *Front. Mar. Sci.* 4 (2017) 72, <https://doi.org/10.3389/fmars.2017.00072>.
- [45] MEDIAS Handbook, 2015, Common protocol for the Pan-Mediterranean Acoustic Survey (MEDIAS), Sète, France, March 2015, p. 21. Available from: <http://medias-project.eu/medias/website/handbooks-menu/func-startdown/60/>.
- [46] B.S. Galil, The limit of the sea: the bathyal fauna of the Levantine Sea, *Sci. Mar.* 68 (S3) (2004) 63–72.
- [47] V. Papiol, J.E. Cartes, E. Fanelli, P. Rumolo, Food web structure and seasonality of slope megafauna in the NW Mediterranean elucidated by stable isotopes: relationship with available food sources, *J. Sea Res.* 77 (2013) 53–69.
- [48] E. Fanelli, E. Azzurro, M. Bariche, J.E. Cartes, F. Maynou, Depicting the novel Eastern Mediterranean food web: a stable isotopes study following Lessepsian fish invasion, *Biol. Invasions* 17 (2015) 2163–2178.
- [49] M. Valls, C.J. Sweeting, M.P. Olivar, M.F. de Puelles, C. Pasqual, N.V.C. Polunin, A. Quetglas, Structure and dynamics of food webs in the water column on shelf

- and slope grounds of the western Mediterranean, *J. Mar. Syst.* 138 (2014) 171–181.
- [50] S. Bianchelli, R. Danovaro, Meiofaunal biodiversity in submarine canyons of the Mediterranean Sea: a meta-analysis, *Prog. Oceanogr.* 170 (2019) 69–80.
- [51] A. Pusceddu, S. Bianchelli, J. Martín, P. Puig, A. Palanques, P. Masqué, R. Danovaro, Chronic and intensive bottom trawling impairs deep-sea biodiversity and ecosystem functioning, *Proc. Natl. Acad. Sci.* 111 (2014) 8861–8866.
- [52] R. Danovaro, C. Corinaldesi, E. Rastelli, A. Dell'Anno, Towards a better quantitative assessment of the relevance of deep-sea viruses, Bacteria and Archaea in the functioning of the ocean seafloor, *Aquat. Microb. Ecol.* 75 (2015) 81–90, <https://doi.org/10.3354/ame01747>.
- [53] F. Semprucci, C. Sbrocca, M. Rocchi, M. Balsamo, Temporal changes of the meiofaunal assemblage as a tool for the assessment of the ecological quality status, *J. Mar. Biol. Assoc. U. K.* 95 (2014) 247–254.
- [54] S. Bianchelli, A. Pusceddu, E. Buschi, R. Danovaro, Trophic status and meiofauna biodiversity in the Northern Adriatic Sea: insights for the assessment of good environmental status, *Mar. Environ. Res.* 113 (2016) 18–30.
- [55] R. Danovaro, C. Gambi, A. Dell'Anno, C. Corinaldesi, S. Fraschetti, A. Vanreusel, M. Vincx, A.J. Gooday, Exponential decline of deep-sea ecosystem functioning linked to benthic biodiversity loss, *Curr. Biol.* 18 (2008) 1–8.
- [56] F. Sardá, A. Calafat, M. Flexas, A. Tselepidis, M. Canals, M. Espino, A. Tursi, An introduction to Mediterranean deep-sea biology, *Sci. Mar.* 68 (suppl. 3) (2004) 7–38.
- [57] E. Fanelli, S. Bianchelli, R. Danovaro, Deep-sea mobile megafauna of Mediterranean submarine canyons and open slopes: analysis of spatial and bathymetric gradients, *Prog. Oceanogr.* 168 (2018) 23–24.
- [58] M. Taviani, L. Angeletti, S. Canese, R. Cannas, F. Cardone, A. Cau, A.B. Cau, M. C. Follesa, F. Marchese, P. Montagna, C. Tessarolo, The “Sardinian cold-water coral province” in the context of the Mediterranean coral ecosystems, *Deep-Sea Res. Part II Top. Stud. Oceanogr.* 145 (2017) 61–78.
- [59] G. Chimienti, M. Bo, M. Taviani, F. Mastroianni, Occurrence and biogeography of cold water coral communities in Mediterranean hard and soft bottoms, in: C. Orejas, C. Jiménez (Eds.), *Mediterranean Cold-Water Corals: Past, Present and Future*, Springer International Publishing AG, part of Springer Nature 2019, 2019. Coral Reefs of the World 9.
- [60] A. Freiwald, L. Beuck, A. Rüggeberg, M. Taviani, D. Hebbeln, R/V Meteor M70-1 Participants, The white coral community in the central Mediterranean Sea revealed by ROV surveys, *Oceanography* 22 (1) (2009) 58–74.
- [61] V. Lauria, G. Garofalo, F. Fiorentino, D. Massi, G. Milisenda, S. Piraino, T. Russo, M. Gristina, Species distribution models of two critically endangered deep-sea octocorals reveal fishing impacts on vulnerable marine ecosystems in central Mediterranean Sea, *Sci. Rep.* 7 (2017) 8049.
- [62] G. Vasilis, C.J. Smith, S. Kiparissis, C. Stamouli, C. Dounas, C. Mytilineou, Updating the distribution status of the critically endangered bamboo coral *Isidella elongata* (Esper, 1788) in the deep Eastern Mediterranean Sea, *Regional Stud. Mar. Sci.* 28 (2019) 100610, <https://doi.org/10.1016/j.rsm.2019.100610>.
- [63] M. Taviani, F. Freiwald, H. Zibrowius, Deep coral growth in the Mediterranean Sea: an overview, in: A. Freiwald, J.M. Roberts (Eds.), *Cold-water Corals and Ecosystems*, Springer-Verlag, Berlin, 2005, pp. 137–156.
- [64] M. Taviani, L. Angeletti, B. Antolini, A. Ceregato, C. Frogli, M. Lopez Correa, P. Montagna, A. Remia, F. Trincardi, A. Vertino, Geo-biology of Mediterranean deep-water coral ecosystems, *Mar. Res. CNR* 6 (2011) 705–719.
- [65] M. Taviani, L. Angeletti, F. Cardone, P. Montagna, R. Danovaro, A unique and threatened deep water coral-bivalve biotope new to the Mediterranean Sea offshore the Naples megalopolis, *Sci. Rep.* 9 (2019) 3411, <https://doi.org/10.1038/s41598-019-39655-8>.
- [66] P.J. Schembri, M. Dimech, M. Camilleri, R. Page, Living deep-water *Lophelia* and *Madrepora* corals in Maltese waters (Strait of Sicily, Mediterranean sea), *Cah. Biol. Mar.* 48 (2007) 77–83.
- [67] M.C. Fabri, L. Pedel, L. Beuck, F. Galgani, D. Hebbeln, A. Freiwald, Megafauna of vulnerable marine ecosystems in French Mediterranean submarine canyons: spatial distribution and anthropogenic impacts, *Deep Sea Res.* 104 (Part I) (2014) 184–207.
- [68] M.C. Fabri, A. Bargain, I. Paireaud, L. Pedel, I. Taupier-Letage, Cold-water coral ecosystems in Cassidaigne Canyon: an assessment of their environmental living conditions, *Deep Sea Res.* 137 (Part II) (2017) 436–453.
- [69] M. Bo, G. Bavestrello, M. Angiolillo, L. Calcagnile, S. Canese, R. Cannas, A. Cau, M. D'Elia, F. D'Oriano, M.C. Follesa, G. Quarta, Persistence of pristine deep-sea coral gardens in the Mediterranean Sea (SW Sardinia), *PLoS One* 10 (3) (2015), e0119393.
- [70] J. Evans, R. Aguilar, H. Alvarez, J.A. Borg, S. Garcia, L. Knittweis, P.J. Schembri, Recent evidence that the deep sea around Malta is a biodiversity hotspot, in: 41st Congress of the Mediterranean Science Commission (CIESM), Kiel, Germany, 12–16 September 2016, 2016.
- [71] E. Fanelli, I. Delbono, R. Ivaldi, M. Pratellesi, S. Cocito, A. Peirano, Cold water coral *Madrepora oculata* in the eastern Ligurian Sea (NW Mediterranean): historical banks and recent findings, *Aquat. Conserv. Mar. Freshw. Ecosyst.* 27 (2017) 965–975.
- [72] D. Moccia, A. Cau, A. Alvito, S. Canese, R. Cannas, M. Bo, M. Angiolillo, M. C. Follesa, New sites expanding the “Sardinian cold-water coral province” extension: a new potential cold-water coral network? *Aquat. Conserv. Mar. Freshw. Ecosyst.* 29 (2019) 153–160.
- [73] S. Migeon, J. Mascle, M. Coste, P. Rouillard, Mediterranean submarine canyons and channels: morphological and geological backgrounds, in: M. Wurtz (Ed.), *Mediterranean Submarine Canyons: Ecology and Governance*, IUCN, Gland, Switzerland and Malaga, Spain, 2012, pp. 27–41.
- [74] M. Wurtz, M. Rovere, Atlas of the Mediterranean Seamounts and Seamount-like Structures, IUCN, Gland, Switzerland and Malaga, Spain, 2015, p. 276.
- [75] J. Mascle, F. Mary, D. Praeg, L. Brosolo, L. Camera, S. Ceramicola, S. Dupré, Distribution and geological control of mud volcanoes and other fluid/free gas seepage features in the Mediterranean Sea and nearby Gulf of Cadiz, *Geo Mar. Lett.* 34 (2014) 89–110.
- [76] R. Koppelman, R. Bottger-Schnack, J. Mobius, H. Weikert, Trophic relationships of zooplankton in the eastern Mediterranean based on stable isotope measurements, *J. Plankton Res.* 31 (2009) 669–686.
- [77] E. Fanelli, V. Papiol, J.E. Cartes, P. Rumolo, C. Brunet, M. Sprovieri, Food web structure of the megabenthic, invertebrate epifauna on the Catalan slope (NW Mediterranean): evidence from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis, *Deep Sea Res., Part I* 58 (2011) 98–109.
- [78] E. Fanelli, J.E. Cartes, V. Papiol, Food web structure of deep-sea macrozooplankton and micronekton off the Catalan slope: insight from stable isotopes, *J. Mar. Syst.* 87 (2011) 79–89.
- [79] E. Fanelli, V. Papiol, J.E. Cartes, O. Rodriguez-Romeu, Trophic ecology of *Lampanyctus crocodilus* on north-west Mediterranean Sea slopes in relation to reproductive cycle and environmental variables, *J. Fish Biol.* 84 (2014) 1654–1688.
- [80] J.E. Cartes, C. Lolocono, V. Mamouridis, C. López-Pérez, P. Rodríguez, Geomorphological, trophic and human influences on the bamboo coral *Isidella elongata* assemblages in the deep Mediterranean: to what extent does *Isidella* form habitat for fish and invertebrates? *Deep Sea Res. Part I* 76 (2013) 52–65.
- [81] A. Denda, B. Christiansen, Zooplankton distribution patterns at two seamounts in the subtropical and tropical NE Atlantic, *Mar. Ecol. Prog. Ser.* 35 (2) (2014) 159–179.
- [82] I. Conese, E. Fanelli, S. Miserocchi, L. Langone, Food web structure and trophodynamics of deep-sea plankton from the Bari Canyon and adjacent slope (Southern Adriatic, central Mediterranean Sea), *Prog. Oceanogr.* 175 (2019) 92–104.
- [83] B.S. Galil, A. Marchini, A. Occhipinti-Ambrogi, D. Minchin, A. Narcius, H. Ojaveer, S. Olenin, International arrivals: widespread bioinvasions in European Seas, *Ethol. Ecol. Evol.* 26 (2–3) (2014) 152–171, <https://doi.org/10.1080/03949370.2014.897651>.
- [84] A. Zenetos, Mediterranean Sea: 30 years of biological invasions (1988–2017), in: 1st Mediterranean Symposium on the Non-Indigenous Species, 2018, 2019, p. 13.
- [85] B.S. Galil, Loss or gain? Invasive aliens and biodiversity in the Mediterranean sea, *Mar. Pollut. Bull.* 55 (7–9) (2007) 314–322, <https://doi.org/10.1016/j.marpolbul.2006.11.008>.
- [86] S. Katsanevakis, G. Verriopoulos, A. Nicolaidou, M. Thessalou-Legaki, Effect of marine litter on the benthic megafauna of coastal soft bottoms: a manipulative field experiment, *Mar. Pollut. Bull.* 54 (2007) 771–778.
- [87] B.S. Galil, A. Marchini, A. Occhipinti-Ambrogi, East is east and West is west? Management of marine bioinvasions in the Mediterranean Sea, *Estuarine, Coastal Shelf Sci.* (2016), <https://doi.org/10.1016/j.ecss.2015.12.021>.
- [88] B. Galil, A. Marchini, A. Occhipinti-Ambrogi, H. Ojaveer, The enlargement of the Suez Canal - Erythraean introductions and management challenges, *Manag. Biol. Invasions* 8 (2) (2017) 141–152, <https://doi.org/10.3391/mbi.2017.8.2.02>.
- [89] M. Goren, B.S. Galil, A. Diamant, N. Stern, Y. Levitt-Barmats, Invading up the food web? Invasive fish in the southeastern Mediterranean Sea, *Mar. Biol.* 163 (2016) 8, <https://doi.org/10.1007/s00227-016-2950-7>.
- [90] E. Azzurro, V. Sbragaglia, J. Cerri, M. Bariche, L. Bolognini, J. Ben Souissi, G. Busoni, S. Coco, A. Chryssanthi, E. Fanelli, R. Ghanem, J. Garrabou, F. Gianni, F. Grati, J. Kollitari, G. Letterio, L. Lipej, C. Mazzoldi, N. Milone, F. Pannacciulli, A. Pešić, Y. Samuel-Rhoads, L. Saponari, J. Tomanic, N.E. Topçu, G. Vargiu, P. Moschella, Climate change, biological invasions, and the shifting distribution of Mediterranean fishes: a large-scale survey based on local ecological knowledge, *Glob. Chang. Biol.* 25 (2019) 2779–2792.
- [91] T. Özcan, A.S. Ateş, T. Katağan, Expanding distribution and occurrence of the indo-pacific stomatopod, *erugosquilla massavensis* (Kossmann, 1880) on the aegean coast of Turkey, *Mediterr. Mar. Sci.* 9 (2) (2008) 115–118.
- [92] M. Corsini-Foka, A. Pancucci-Papadopoulou, G. Kondilatos, S. Kalogirou, *Gonioinfradens paucidentatus* (A. Milne Edwards, 1861) (Crustacea, Decapoda, Portunidae): a new alien crab in the Mediterranean sea, *Mediterr. Mar. Sci.* 11 (2010) 331–340.
- [93] G. Innocenti, G. Stasolla, M. Goren, N. Stern, Y. Levitt-Barmats, A. Diamant, B. S. Galil, Going down together: invasive host, *Charybdis longicollis* (Leene, 1938) (Decapoda: Brachyura: Portunidae) and invasive parasite, *Heterosaccus dollfusii* Boschma, 1960 (Cirripedia: Rhizocephala: Sacculinidae) on the upper slope off the Mediterranean coast of Israel, *Mar. Biol. Res.* 13 (2017) 229–236.
- [94] E. Özgür Özbek, M. Cardak, T. Kebapçioğlu, Spatio-temporal patterns of abundance, biomass and length of the silver-cheeked toadfish *Lagocephalus scleratus* in the Gulf of Antalya, *Turk. J. Fish. Aquat. Sci.* 17 (2017) 725–733.
- [95] D. Yağlıoğlu, D. Ayas, New occurrence data of four alien fishes (*Pisodonophis semicinctus*, *Pterois miles*, *Scarus ghobban* and *Parupeneus forsskali*) from the north eastern Mediterranean (Yeşilovacik Bay, Turkey), *Biharean Biologist* 10 (2016) 150–152.
- [96] C. Jimenez, P. Patsalou, V. Andreou, M.F. Huseyinoglu, B.A. Çiçek, L. Hadjiannou, A. Petrou, Out of sight, out of reach, out of mind: invasive lionfish *Pterois miles* in Cyprus at depths beyond recreational diving limits, in: 1st Mediterranean Symposium on the Non-Indigenous Species (Antalya, Turkey, 17–18 January 2019), 2019, pp. 59–64.

- [97] B.S. Galil, R. Danovaro, S.B.S. Rothman, R. Gevili, M. Goren, Invasive biota in the deep-sea Mediterranean: an emerging issue in marine conservation and management, *Biol. Invasions* 21 (2) (2018) 281–288.
- [98] V. Parravicini, E. Azzurro, M. Kulbicki, J. Belmaker, Niche shift can impair the ability to predict invasion risk in the marine realm: an illustration using Mediterranean fish invaders, *Ecol. Lett.* 18 (2015) 246–253.
- [99] D. Gascuel, M. Coll, C. Fox, S. Guénette, J. Guitton, A. Kenny, L. Knittweis, J. R. Nielsen, G. Piet, T. Raid, M. Travers-Trolet, S. Shephard, Fishing impact and environmental status in European seas: a diagnosis from stock assessments and ecosystem indicators, *Fish Fish.* 17 (2016) 31–55.
- [100] M.L. Pinsky, S.R. Palumbi, Meta-analysis reveals lower genetic diversity in overfished populations, *Mol. Ecol.* 23 (1) (2014) 29–39.
- [101] F.W. Allendorf, O. Berry, N. Ryman, So long to genetic diversity, and thanks for all the fish, *Mol. Ecol.* 23 (1) (2014) 23–25.
- [102] E. Fanelli, F. Badalamenti, G. D'Anna, P. Pipitone, C. Romano, Trophodynamic effects of trawling on the feeding ecology of Pandora, *Pagellus erythrinus* off the northern Sicily coast (Mediterranean Sea), *Mar. Freshw. Res.* 61 (2010) 408–417.
- [103] F. Colloca, M. Cardinale, F. Maynou, M. Giannoulaki, G. Scarcella, K. Jenko, J. M. Bellido, F. Fiorentino, Rebuilding Mediterranean fisheries: a new paradigm for ecological sustainability, *Fish Fish.* 14 (2013) 89–109, <https://doi.org/10.1111/j.1467-2979.2011.00453.x>.
- [104] FAO, in: Report of the FAO Workshop on Deep-Sea Fisheries and Vulnerable Marine Ecosystems of the Mediterranean, Rome, Italy, 18–20 July 2016, FAO Fisheries and Aquaculture, Rome, Italy, 2016. Report No. 1183.
- [105] GFCM, in: Report of the First Meeting of the Working Group on Vulnerable Marine Ecosystems (WGVME), Malaga, Spain, 3–5 April 2017, 2017.
- [106] FAO, International Guidelines for the Management of Deep-Sea Fisheries in the High Seas, FAO, Rome, Italy, 2009, p. 73pp.
- [107] FAO, The State of Mediterranean and Black Sea Fisheries, General Fisheries Commission for the Mediterranean, Rome, 2018, p. 172pp.
- [108] G. Le Corre, H. Farrugio, Note sur la création par la CGPM d'une Zone de pêche réglementée dans le golfe du Lion en mars 2009, 2011. RBE/HMT 2011-002, <https://archimer.ifremer.fr/doc/00086/19688/>.
- [109] R. Elahi, F. Ferretti, A. Bastari, C. Cerrano, F. Colloca, J. Kowalik, M. Ruckelshaus, A. Struck, F. Micheli, Leveraging vessel traffic data and a temporary fishing closure to inform marine management, *Front. Ecol. Environ.* 16 (8) (2018) 1–7.
- [110] ICES, EU request to provide guidance on operational methods for the evaluation of the MSFD Criterion D3C3, in: Report of the ICES Advisory Committee, 2016. ICES Advice 2016, Book 1, Section 1.6.2.2, 2016.
- [111] ICES, Report of the Workshop on Guidance on Development of Operational Methods for the Evaluation of the MSFD Criterion D3.3 (WKIND3.3ii), 1–4 November 2016, vol. 44, ICES, Copenhagen, Denmark, 2017, p. 145. CM 2016/ACOM.
- [112] L. Thomsen, J. Aguzzi, C. Costa, F. De Leo, A. Ogston, A. Purser, The oceanic biological pump: rapid carbon transfer to the Deep Sea during winter, *Sci. Rep.* 7 (2017) 10763.
- [113] D. van Oevelen, M. Bergmann, K. Soetaert, E. Bauerfeind, C. Hasemann, M. Klages, I. Schewe, T. Soltwedel, N.E. Budaeva, Carbon flows in the benthic food web at the deep-sea observatory HAUSGARTEN (Fram Strait), *Deep Sea Res., Part I* 58 (2011) 1069–1083.
- [114] C. Gambi, A. Pusceddu, L. Benedetti-Cecchi, R. Danovaro, Species richness, species turnover and functional diversity in nematodes of the deep Mediterranean sea: searching for drivers at different spatial scales, *Glob. Ecol. Biogeogr.* 23 (1) (2014) 24–39.
- [115] ICES, Report of the Workshop on Guidance for the Review of MSFD Decision Descriptor 3 - Commercial Fish and Shellfish II (WKGMSFDD3-II), 10–12 February 2015, ICES Headquarters, Denmark, vol. 48, ICES, 2015, p. 36. CM 2015/ACOM.
- [116] I. Rombouts, G. Beaugrand, L.F. Artigas, J.C. Dauvin, F. Gevaert, E. Goberville, D. Kopp, S. Lefebvre, C. Luczak, N. Spilmont, M. Travers-Trolet, M.C. Villanueva, R.R. Kirby, Evaluating marine ecosystem health: case studies of indicators using direct observations and modelling methods, *Ecol. Indic.* 24 (2013) 353–365.
- [117] E. Fanelli, J.E. Cartes, P. Rumolo, M. Sprovieri, Food-web structure and trophodynamics of mesopelagic-suprabenthic bathyal macrofauna of the Algerian Basin based on stable isotopes of carbon and nitrogen, *Deep Sea Res., Part I* 56 (2009) 1504–1520.
- [118] K. Iken, T. Brey, U. Wand, J. Voigt, P. Junghans, Food web structure of the benthic community at the Porcupine Abyssal Plain (NE Atlantic): a stable isotope analysis, *Prog. Oceanogr.* 50 (1–4) (2001) 383–405.
- [119] MEDITS Handbook. Version N. 9, MEDITS Working Group, 2017, p. 106. Available from: <http://www.sibm.it/MEDITS%202011/principaledownload.htm>.
- [120] C. Stefanescu, B. Moralesnin, E. Massuti, Fish assemblages on the slope in the Catalan sea (western Mediterranean) - influence of a submarine-canyon, *J. Mar. Biol. Assoc. U. K.* 74 (1994) 499–512.
- [121] E. Massuti, J.D. Gordon, J. Moranta, S.C. Swan, C. Stefanescu, Mediterranean and Atlantic deep-sea fish assemblages: differences in biomass composition and size-related structure, *Sci. Mar.* 68 (S3) (2004) 101–115.
- [122] A. Anastasopoulou, F. Biandolino, A. Chatzisyrou, F. Hemida, B. Guijarro, V. Kousteni, Ch Mytilineou, P. Pattoura, E. Prato, New fisheries-related data from the Mediterranean sea (november, 2016), *Mediterr. Mar. Sci.* 17 (2016) 822–827.
- [123] E. Fanelli, J.E. Cartes, V. Papiol, C. López-Pérez, M. Carrasson, Long-term decline in the trophic level of megafauna in the deep Mediterranean Sea: a stable isotopes approach, *Clim. Res.* 67 (2016) 191–207.
- [124] P. Cresson, M.C. Fabri, M. Bouchouca, C. Brach Papa, F. Chavanon, A. Jadaud, J. Knoery, F. Miralles, D. Cossa, Mercury in organisms from the Northwestern Mediterranean slope: importance of food sources, *Sci. Total Environ.* 497–498 (2014), 229–23.
- [125] A. Carlier, E. Le Guilloux, K. Olu, J. Sarrazin, F. Mastrototaro, M. Taviani, J. Clavier, Trophic relationships in a deep mediterranean cold-water coral bank (santa maria di Leuca, Ionian Sea), *Mar. Ecol. Prog. Ser.* 397 (2009) 125–137.
- [126] S. Tecchio, D. van Oevelen, K. Soetaert, J. Navarro, E. Ramirez-Llodra, Trophic dynamics of deep-sea megabenthos are mediated by surface productivity, *PLoS One* 8 (5) (2013), e63796, <https://doi.org/10.1371/journal.pone.0063796>.
- [127] J.E. Cartes, E. Fanelli, K. Kaporis, Y.K. Bayhan, A. Ligas, C. López-Pérez, M. Murenu, V. Papiol, P. Rumolo, G. Scarcella, Spatial variability in the trophic ecology and biology of the deep-sea shrimp *Aristaeomorpha foliacea* in the Mediterranean Sea, *Deep Sea Res. Part I* 87 (Supplement C) (2014) 1–13.
- [128] M.S. Naumann, I. Imma Tolosa, M. Taviani, R. Grover, C. Christine Ferrier-Pagés, Trophic ecology of two cold-water coral species from the Mediterranean Sea revealed by lipid biomarkers and compound-specific isotope analyses, *Coral Reefs* 34 (2015) 1165–1175, <https://doi.org/10.1007/s00338-015-1325-8>.
- [129] G.M. Luna, S. Bianchelli, F. Decembrini, E. De Domenico, R. Danovaro, A. Dell'Anno, The dark portion of the Mediterranean Sea is a bioreactor of organic matter cycling, *Glob. Biogeochem. Cycles* 26 (2) (2012) GB2017, <https://doi.org/10.1029/2011GB004168>.
- [130] M. Molari, E. Manini, A. Dell'Anno, Dark inorganic carbon fixation sustains the functioning of benthic deep-sea ecosystems, *Glob. Biogeochem. Cycles* 27 (2013) 212–221.
- [131] J.G. Ferreira, J.H. Andersen, A. Borja, S.B. Bricker, J. Camp, M. Cardoso da Silva, E. Garcés, A.S. Heiskanen, C. Humborg, L. Ignatiades, C. Lancelot, A. Menesguen, P. Tett, N. Hoepffner, U. Claussen, Marine Strategy Framework Directive Task Group 5 Report Eutrophication, 2010. EUR 24338.
- [132] J.G. Ferreira, C. Vale, C.V. Soares, F. Salas, P.E. Stacey, S.B. Bricker, M.C. Silva, J. C. Marques, Monitoring of coastal and transitional waters under the EU water framework directive, *Environ. Monit. Assess.* 135 (2007) 195–216.
- [133] Y.J. Xiao, J.G. Ferreira, S.B. Bricker, J.P. Nunes, M.Y. Zhu, X.L. Zhang, Trophic assessment in Chinese coastal systems - review of methods and application to the Changjiang (Yangtze) estuary and Jiaozhou Bay, *Estuar. Coasts* 30 (2007) 901–918.
- [134] A. Borja, S.B. Bricker, D.M. Daur, N.T. Demetriades, J.G. Ferreira, A.T. Forbes, P. Hutchings, X.P. Jia, R. Kenchington, J.C. Marques, et al., Overview of integrative tools and methods in assessing ecological integrity in estuarine and coastal systems worldwide, *Mar. Pollut. Bull.* 56 (9) (2008) 1519–1537.
- [135] S.B. Bricker, B. Longstaff, W. Dennison, A. Jones, K. Boicourt, C. Wicks, J. Woerner, Effects of nutrient enrichment in the nation's estuaries: a decade of change, *Harmful Algae* 8 (1) (2008) 21–32.
- [136] S.W. Nixon, Eutrophication and the microscope, in: *Eutrophication in Coastal Ecosystems*, Springer, Dordrecht, 2009, pp. 5–19.
- [137] U. Claussen, W. Zevenboom, U. Brockmann, D. Topcu, P. Bot, Assessment of the eutrophication status of transitional, coastal and marine waters within OSPAR, *Hydrobiologia* 629 (1) (2009) 49–58, <https://doi.org/10.1007/s10750-009-9763-3>.
- [138] J.G. Ferreira, J.H. Andersen, A. Borja, S.B. Bricker, J. Camp, M. Cardoso da Silva, E. Garcés, A.S. Heiskanen, C. Humborg, L. Ignatiades, C. Lancelot, A. Menesguen, P. Tett, N. Hoepffner, U. Claussen, Overview of eutrophication indicators to assess environmental status within the European Marine Strategy Framework Directive, *Estuar. Coast Shelf Sci.* 93 (2011) 117–131.
- [139] C. Doya, J. Aguzzi, D. Chatziveangelou, C. Costa, J.B. Company, V. Tunnicliffe, The seasonal use of small-scale space by benthic species in a transiently hypoxic area, *J. Mar. Syst.* 154 (2015) 280–290.
- [140] D. Breitburg, L.A. Levin, A. Oschlies, M. Grégoire, F.P. Chavez, D.J. Conley, V. Garçon, D. Gilbert, D. Gutiérrez, K. Isensee, G.S. Jacinto, Declining oxygen in the global ocean and coastal waters, *Science* 359 (6371) (2018).
- [141] R.F. Keeling, A. Kortzinger, N. Gruber, Ocean deoxygenation in a warming world, *Ann. Rev. Mar. Sci.* 2 (2010) 199–229, <https://doi.org/10.1146/annurev.marine.010908.163855>.
- [142] L. Stramma, G.C. Johnson, J. Sprintall, V. Mohrholz, Expanding oxygen-minimum zones in the tropical oceans, *Science* 320 (5876) (2008) 655–658.
- [143] L.A. Levin, Oxygen minimum zone benthos: adaptation and community response to hypoxia, *Oceanogr. Mar. Biol.: Ann. Rev.* 2003 41 (2003) 1–45.
- [144] R. Danovaro, P.V.R. Snelgrove, P. Tyler, Challenging the paradigms of deep-sea ecology, *Trends Ecol. Evol.* 29 (8) (2014) 465–475.
- [145] C. Piroddi, M. Coll, C. Lique, D. Macias, K. Greer, J. Buszewski, J. Steenbeek, R. Danovaro, V. Christensen, Historical changes of the Mediterranean Sea ecosystem: modelling the role and impact of primary productivity and fisheries changes over time, *Sci. Rep.* 7 (2017) 44491, <https://doi.org/10.1038/srep44491>.
- [146] E. Ramirez-Llodra, P.A. Tyler, M.C. Baker, O.A. Bergstad, M.R. Clark, E. Escobar, L.A. Levin, L. Menot, A.A. Rowden, C.R. Smith, C.L. van Dover, Man and the last great wilderness: human impact on the deep sea, *PLoS One* 6 (8) (2011), e22588.
- [147] A. Dell'Anno, M.L. Mei, A. Pusceddu, R. Danovaro, Assessing the trophic state and eutrophication of coastal marine systems: a new approach based on the biochemical composition of sediment organic matter, *Mar. Pollut. Bull.* 44 (2002) 611–622.
- [148] M.C. Fabri, A. Brind'Amour, A. Jadaud, F. Galgani, S. Vaz, M. Taviani, G. Scarcella, M. Canals, A. Sanchez, J. Grimalt, B. Galil, M. Goren, P. Schembri, J. Evans, Leyla Knittweis, A.-L. Cantafaro, E. Fanelli, L. Carugati, R. Danovaro, Review of Literature on the Implementation of the MSFD to the Deep Mediterranean Sea. IDEM Project, 2018, p. 228 p, <https://doi.org/10.13155/53809.Deliverable.1.1>, www.msfd-idem.eu.
- [149] J.M. Mercado, L. Yebra, D. Cortés, C. Beken, M. Simbora, S. Moncheva, A. Alonso, F. Gómez, S. Salles, A. Sánchez, N. Valcarcel, Designing joint monitoring programs for the MSFD Eutrophication assessment based on the

- monitoring strategy of UNEP/MAP (Barcelona Convention), in: *Plans for the Design of Joint Monitoring Programs in the Mediterranean and Black Sea Regions Adapted to MSFD Requirements*. - IRIS-SES Project. Fransisco Alemany, Pagou Kalliopi, Giannoudi Louisa, Streftaris Nikos, IRIS-SES Project, 2015. May 2015.
- [150] A. Pusccheddu, A. Dell'Anno, M. Fabiano, R. Danovaro, Quantity and bioavailability of sediment organic matter as signature of benthic trophic state, *Mar. Ecol. Prog. Ser.* 375 (2009) 41–52.
- [151] UNEP(DEPI)/MED, Eutrophication Monitoring Strategy for the MED POL (REVISION) UNEP/MAP; WG.321/Inf.5, 9 November 2007, 2007.
- [152] A. Pusccheddu, S. Bianchelli, C. Gambi, R. Danovaro, Assessment of benthic trophic status of marine coastal ecosystems: significance of meiofaunal rare taxa, *Estuar. Coast Shelf Sci.* 93 (2011) 420–430.
- [153] S. Bianchelli, E. Buschi, R. Danovaro, A. Pusccheddu, Biodiversity loss and turnover in alternative states in the Mediterranean Sea: a case study on meiofauna, *Sci. Rep.* 6 (2016) 34544.
- [154] R.E. Boschen, A.A. Rowden, M.R. Clark, J.P.A. Gardner, Mining of deep-sea seafloor massive sulfides: a review of the deposits, their benthic communities, impacts from mining, regulatory frameworks and management strategies, *Ocean Coast Manag.* 84 (Supplement C) (2013) 54–67.
- [155] P. Holler, E. Markert, A. Bartholoma, R. Capperucci, H.C. Hass, I. Kroncke, F. Mielck, H.C. Reimers, Tools to evaluate seafloor integrity: comparison of multi-device acoustic seafloor classifications for benthic macrofauna-driven patterns in the German Bight, southern North Sea, *Geo Mar. Lett.* 37 (2017) 93–109.
- [156] P. Puig, M. Canals, J.B. Company, J. Martin, D. Amblas, G. Lastras, A. Palanques, A.M. Calafat, Ploughing the deep sea floor, *Nature* 489 (2012) 286–289, <https://doi.org/10.1038/nature11410>, 7415.
- [157] J. Martin, P. Puig, P. Masque, A. Palanques, A. Sanchez-Gomez, Impact of bottom trawling on deep-sea sediment properties along the flanks of a submarine canyon, *PLoS One* 9 (2014), e104536, <https://doi.org/10.1371/journal.pone.0104536>, 8.
- [158] O.R. Eigaard, F. Bastardie, N.T. Hintzen, L. Buhl-Mortensen, P. Buhl-Mortensen, R. Catarino, G.E. Dinesen, J. Egekvist, H.O. Fock, K. Geitner, H.D. Gerritsen, M. M. González, P. Jonsson, S. Kavadas, P. Laffargue, M. Lundy, G. Gonzalez-Mirelis, J.R. Nielsen, N. Papadopoulou, E. Posen, P.E. J. Pulcinella, T. Russo, A. Sala, C. Silva, C.J. Smith, B. Vanelslander, A.D. Rijnsdorp, The footprint of bottom trawling in European waters: distribution, intensity, and seabed integrity, *ICES J. Mar. Sci.* 74 (2017) 847–865.
- [159] J. Martin, P. Puig, A. Palanques, P. Masque, J. Garcia-Orellana, Effect of commercial trawling on the deep sedimentation in a Mediterranean submarine canyon, *Mar. Geol.* 252 (2008) 150–155.
- [160] The Petroleum Economist Ltd, World Energy Atlas, seventh ed., SC (Sang Choy). International Pte Ltd, Singapore, 2013. ISBN 1 86186 343 8.
- [161] B. Galil, B. Herut, Marine Environmental Issues of Deep-Sea Oil and Gas Exploration and Exploitation Activities off the Coast of Israel, 2011, p. 24. IOLR Report H15/2011.
- [162] M. Livnat, Offshore Safety in the Eastern Mediterranean Energy Sector. Implications of the New EU Directive, Mediterranean Paper Series 2014, The German Marshall Fund of the United States, Washington DC, USA, 2014, p. 12.
- [163] J.C. Dauvin, Towards an impact assessment of bauxite red mud waste on the knowledge of the structure and functions of bathyal ecosystems: the example of the Cassidaigne canyon (north-western Mediterranean Sea), *Mar. Pollut. Bull.* 60 (2010) 197–206.
- [164] C. Fontanier, M.C. Fabri, R. Buscail, L. Biscara, K. Koho, G.J. Reichart, D. Cossa, S. Galaup, G. Chabaud, L. Pigot, Deep-sea foraminifera from the Cassidaigne Canyon (NW Mediterranean): assessing the environmental impact of bauxite red mud disposal, *Mar. Pollut. Bull.* 64 (2012) 1895–1910, <https://doi.org/10.1016/j.marpolbul.2012.06.016>.
- [165] C. Fontanier, L. Biscara, B. Mamo, E. Delord, Deep-sea benthic foraminifera in an area around the Cassidaigne Canyon (NW Mediterranean) affected by bauxite discharges, *Mar. Biodivers.* 44 (2014) 4, <https://doi.org/10.1007/s12526-014-0281-9>.
- [166] S. Varnavas, G. Ferentinos, M. Collins, Dispersion of bauxitic red mud in the gulf-of-Corinth, Greece, *Mar. Geol.* 70 (3–4) (1986) 211–222, [https://doi.org/10.1016/0025-3227\(86\)90003-4](https://doi.org/10.1016/0025-3227(86)90003-4).
- [167] S.P. Varnavas, P.P. Achilleopoulos, Factors controlling the vertical and spatial transport of metal-rich particulate matter in seawater at the outfall of bauxitic red mud toxic waste, *Sci. Total Environ.* 175 (1995) 199–205.
- [168] S.E. Poulos, M.B. Collins, C. Pattiaratchi, A. Cramp, W. Gull, M. Tsimplis, G. Papatheodorou, Oceanography and sedimentation in the semi-enclosed, deep-water Gulf of Corinth (Greece), *Mar. Geol.* 134 (1996) 213–235.
- [169] N. Kress, G. Fainshtein, H. Hornung, MAP Technical Reports Series, Monitoring of Hg, Cd, Cu, Pb, Zn, Co, Be and V at a Deep Water Coal Fly Ash Dumping Site, vol. 104, UNEP/FAO, 1996.
- [170] N. Kress, H. Hornung, B. Herut, Concentrations of Hg, Cd, Cu, Zn, Fe and Mn in deep sea benthic fauna from the southeastern Mediterranean Sea: a comparison study between fauna collected at a pristine area and at two waste disposal sites, *Mar. Pollut. Bull.* 36 (1998) 911–921.
- [171] B. Herut, B. Galil, E. Shefer, Monitoring Alpha – disposal site of dredged material in the Mediterranean Sea 9 (2010) 47. Results of monitoring July–September 2009. Israel Oceanographic & Limnological Research Reports (in Hebrew).
- [172] A.J. Kenny, I. Cato, M. Desprez, G. Fader, R.T.E. Schuttenhelm, J. Side, An overview of seabed-mapping technologies in the context of marine habitat classification, *ICES J. Mar. Sci.* 60 (2) (2003) 411–418.
- [173] C.J. Brown, S.J. Smith, P. Lawton, J.T. Anderson, Benthic habitat mapping: a review of progress towards improved understanding of the spatial ecology of the seafloor using acoustic techniques, *Estuar. Coast Shelf Sci.* 92 (2011) 502–520.
- [174] D. Stephens, M. Diesing, A comparison of supervised classification methods for the prediction of substrate type using Multibeam acoustic and legacy grain-size data, *PLoS One* 9 4 (2014), <https://doi.org/10.1371/journal.pone.0093950> e93950.
- [175] G. Lastras, M. Canals, E. Ballesteros, J.M. Gili, A. Sanchez-Vidal, Cold-water corals and anthropogenic impacts in La Fonera submarine canyon head, *PLoS One* 16 (5) (2016), e0155729, 11.
- [176] C.J. Brown, P. Blondel, Developments in the application of multibeam sonar backscatter for seafloor habitat mapping, *Appl. Acoust.* 70 (2009) 1242–1247.
- [177] C. Lo Iacono, E. Gràcia, R. Bartolomé, E. Coiras, J.J. Danobeitia, J. Acosta, The habitats of the chella bank, eastern Alboran Sea (western mediterranean), in: P. Harris, E. Baker (Eds.), *Seafloor Geomorphology as Benthic Habitat*: GeoHab Atlas of Seafloor Geomorphic Features and Benthic Habitats, Elsevier, London, 2012, pp. 681–687.
- [178] A. Bargain, F. Marchese, A. Savini, M. Taviani, M.-C. Fabri, Santa Maria di Leuca Province (Mediterranean Sea): identification of suitable mounds for cold-water coral settlement using geomorphometric proxies and Maxent methods, *Front. Mar. Sci.* 4 (2017) 338, <https://doi.org/10.3389/fmars.2017.00338>.
- [179] A. Bargain, F. Fogliini, I. Pairaud, D. Bonaldo, S. Carniel, L. Angeletti, M. Taviani, S. Rochette, M.-C. Fabri, Predictive habitat modeling in two Mediterranean canyons including hydrodynamics variables, *Prog. Oceanogr.* 169 (2018) 151–168.
- [180] L. Angeletti, A. Bargain, E. Campiani, F. Fogliini, V. Grande, Leidi, A. Mercorella, M. Prampolini, M. Taviani, Cold-water coral habitat mapping in the Mediterranean sea: methodologies and perspectives, in: C. Orejas, C. Jiménez (Eds.), *Mediterranean Cold-Water Corals: Past, Present and Future*, Springer International Publishing AG, part of Springer Nature, 2019, https://doi.org/10.1007/978-3-319-91608-8_16. Coral Reefs of the World 9.
- [181] C. Lo Iacono, K. Robert, R. Gonzalez-Villanueva, A. Gori, J.-M. Gili, C. Orejas, Predicting cold-water coral distribution in the Cap de Creus Canyon (NW Mediterranean): implications for marine conservation planning, *Prog. Oceanogr.* 169 (2018) 169–180.
- [182] M. Giusti, C. Innocenti, S. Canese, Predicting suitable habitat for the gold coral savaglia savaglia (Bertoloni, 1819) (Cnidaria, Zoantharia) in the south Tyrrhenian Sea, *Cont. Shelf Res.* 81 (2014) 19–28, <https://doi.org/10.1016/j.csr.2014.03.011>.
- [183] M. Giusti, M. Bo, M. Angiolillo, R. Cannas, A. Cau, M.C. Follesa, S. Canese, Habitat preference of *Viminella flagellum* (Alcyonacea: ellisellidae) in relation to bathymetric variables in southeastern Sardinian waters, *Cont. Shelf Res.* 138 (2017) 41–50, <https://doi.org/10.1016/j.csr.2017.03.004>.
- [184] K. Schroeder, G.P. Gasparini, M. Borghini, A. Ribotti, Experimental evidences of the recent abrupt changes in the deep Western Mediterranean Sea, *Dyn. Mediterranean Deep Waters* 38 (2009) (Qwara, Malta).
- [185] K. Schroeder, J. Chiggiato, H.L. Bryden, M. Borghini, S. Ben Ismail, Abrupt climate shift in the western Mediterranean sea, *Sci. Rep.* 6 (2016) 23009, <https://doi.org/10.1038/srep23009>.
- [186] P.C. Reid, R.E. Hari, G. Beaugrand, D.M. Livingstone, C. Marty, D. Straile, J. Barichivich, E. Goberville, R. Adrian, Y. Aono, R. Brown, J. Foster, P. Groisman, P. Héliouët, H. Hsu, R. Kirby, J. Knight, A. Kraberg, J. Li, T. Lo, R.B. Myneni, R. P. North, J.A. Pounds, T. Sparks, R. Stübi, Y. Tian, K.H. Wiltshire, D. Xiao, Z. Zhu, Global impacts of the 1980s regime shift, *Glob. Chang. Biol.* 22 (2) (2016) 682–703.
- [187] A. Conversi, S. Fonda-Umani, T. Peluso, J.C. Molinero, A. Santojanni, M. Edwards, The Mediterranean sea regime shift at the end of the 1980s, and intriguing Parallelisms with other European basins, *PLoS One* 5 (2010) 5, <https://doi.org/10.1371/journal.pone.0010633>.
- [188] W. Roether, R. Well, Oxygen consumption in the eastern mediterranean, *Deep Sea Res., Part 1* 48 (2001) 1535–1551.
- [189] B. Klein, W. Roether, N. Kress, B.B. Manca, M.R. d'Alcala, E. Souvermezoglou, A. Theoharis, G. Civitarese, A. Luchetta, Accelerated oxygen consumption in eastern Mediterranean deep waters following the recent changes in thermohaline circulation, *J. Geophys. Res. Oceans* 108 (2003) C9, <https://doi.org/10.1029/2002jc001454>.
- [190] B. Klein, W. Roether, B.B. Manca, D. Bregant, V. Beitzel, V. Kovacevic, A. Luchetta, The large deep water transient in the Eastern Mediterranean, *Deep Sea Res., Part 1* 46 (1999) 371–414.
- [191] A. Palanques, X.D. de Madron, P. Puig, J. Fabres, J. Guillen, A. Calafat, M. Canals, S. Heussner, J. Bonnin, Suspended sediment fluxes and transport processes in the Gulf of Lions submarine canyons. The role of storms and dense water cascading, *Mar. Geol.* 234 (2006) 43–61.
- [192] J. Martin, J.C. Miquel, A. Khrifpounoff, Impact of open sea deep convection on sediment remobilization in the western Mediterranean, *Geophys. Res. Lett.* 37 (2010), <https://doi.org/10.1029/2010gl043704>.
- [193] A. Sanchez-Vidal, C. Pasqual, P. Kerherve, A. Calafat, S. Heussner, A. Palanques, X. Durrieu de Madron, M. Canals, P. Puig, Impact of dense shelf water cascading on the transfer of organic matter to the deep western Mediterranean basin, *Geophys. Res. Lett.* 35 (2008) L05605, <https://doi.org/10.1029/2007GL032825>.
- [194] J.B. Company, P. Puig, F. Sarda, A. Palanques, M. Latasa, R. Scharek, Climate Influence on deep sea populations, *PLoS One* 3 (2008), e1431, <https://doi.org/10.1371/journal.pone.0001431>, 1.
- [195] M. Rixen, J.M. Beckers, S. Levitus, J. Antonov, T. Boyer, C. Maillard, M. Fichaut, E. Balopoulos, S. Iona, H. Dooley, M.-J. Garcia, B. Manca, A. Giorgetti, G. Manzella, N. Mikhailov, N. Pinardi, M. Zavatarelli, The western mediterranean deep water: a proxy for climate change, *Geophys. Res. Lett.* 32 (2005) 12, <https://doi.org/10.1029/2005gl022702>.

- [196] N. Kress, I. Gertman, B. Herut, Temporal evolution of physical and chemical characteristics of the water column in the Easternmost Levantine basin (Eastern Mediterranean Sea) from 2002 to 2010, *J. Mar. Syst.* 135 (2014) 6–13, <https://doi.org/10.1016/j.jmarsys.2013.11.016>.
- [197] A. Monaco, S. Peruzzi, The Mediterranean Targeted Project MATER—a multiscale approach of the variability of a marine system—overview, *J. Mar. Syst.* 33 (2002) 3–21.
- [198] M. Fichaut, M.J. Garcia, A. Giorgetti, A. Iona, A. Kuznetsov, M. Rixen, M. Group, MEDAR/MEDATLAS 2002: a Mediterranean and Black Sea database for operational oceanography, in: H. Dahlin, N.C. Flemming, K. Nittis, S.E. Petersson (Eds.), Elsevier Oceanography Series, vol. 69, Elsevier, 2003, pp. 645–648, [https://doi.org/10.1016/S0422-9894\(03\)80107-1.0422-9894](https://doi.org/10.1016/S0422-9894(03)80107-1.0422-9894).
- [199] S. Simoncelli, C. Coatanoan, V. Myroshnychenko, H. Sagen, Ö. BÄck, S. Scory, A. Grandi, R. Schlitzer, M. Fichaut, Second release of the SeaDataNet aggregated data sets products, SeaDataNet (2015), <https://doi.org/10.13155/50382>. <http://archimer.ifremer.fr/doc/00392/50382/>.
- [200] C.R. Smith, F.C. De Leo, A.F. Bernardino, A.K. Sweetman, P.M. Arbizu, Abyssal food limitation, ecosystem structure and climate change, *Trends Ecol. Evol.* 23 (2008) 518–528.
- [201] D. Bonaldo, A. Benetazzo, A. Bergamasco, E. Campiani, F. Fogliani, M. Scialvo, F. Trincardi, S. Carniel, Interactions among Adriatic continental margin morphology, deep circulation and bedform patterns, *Mar. Geol.* 375 (2015) 82–98.
- [202] C. Estournel, V. Zervakis, P. Marsaleix, A. Papadopoulos, F. Auclair, L. Perivoliotis, E. Tragou, Dense water formation and cascading in the Gulf of Thermaikos (North Aegean), from observations and modelling, *Cont. Shelf Res.* 25 (2005) 2366–2386.
- [203] K. Schroeder, C. Millot, L. Bengara, S. Ben Ismail, M. Bensi, M. Borghini, G. Budillon, V. Cardin, L. Coppola, C. Curtil, A. Drago, B. El Moumni, J. Font, J. L. Fuda, J. Garcia-Lafuente, G.P. Gasparini, H. Kontoyiannis, D. Lefevre, P. Puig, P. Raimbault, G. Rougier, J. Salat, C. Sammari, J.C. Sanchez Garrido, A. Sanchez-Roman, S. Sparnocchia, C. Tamburini, I. Taupier-Letage, A. Theocharis, M. Vargas-Yanez, A. Vetrano, Long-term monitoring programme of the hydrological variability in the Mediterranean Sea: a first overview of the HYDROCHANGES network, *Ocean Sci.* 9 (2013) 301–324.
- [204] C. Tamburini, M. Canals, X. Durrieu de Madron, L. Houpert, D. Lefevre, S. Martini, ANTARES consortium, deep-sea Bioluminescence blooms after dense water formation at the ocean surface, *PLoS One* 8 (7) (2013), e67523.
- [205] J. Aguzzi, E. Fanelli, T. Ciuffardi, C. Doya, M. Kawato, M. Miyazaki, Y. Furushima, C. Costa, Y. Fujiwhara, Faunal activity rhythms influencing early community succession of an implanted whale carcass offshore Sagami Bay, Japan, *Sci. Rep.* 8 (2018) 11163.
- [206] B.S. Halpern, S. Walbridge, K.A. Selkoe, C.V. Kappel, F. Micheli, C. D'Agrosa, J. F. Bruno, K.S. Casey, C. Ebert, H.E. Fox, R. Fujita, D. Heinemann, H.S. Lenihan, E. M.P. Madin, M.T. Perry, E.R. Selig, M. Spalding, R. Steneck, R. Watson, A global map of human impact on marine ecosystems, *Science* 319 (5865) (2008) 948–952.
- [207] H. Iwata, S. Tanabe, N. Sakai, A. Nishimura, R. Tatsukawa, Geographical distribution of persistent organochlorines in air, water and sediments from Asia and Oceania, and their implications for global redistribution from lower latitudes, *Environ. Pollut.* 85 (1994) 15–33.
- [208] S. Heussner, X. Durrieu de Madron, A. Calafat, M. Canals, J. Carbonne, N. Delsaut, G. Saragoni, Spatial and temporal variability of downward particle fluxes on a continental slope: lessons from an 8-yr experiment in the Gulf of Lions (NW Mediterranean), *Mar. Geol.* 234 (2006) 63–92.
- [209] J. Martin, A. Palanques, P. Puig, Composition and variability of downward particulate matter fluxes in the Palamos submarine canyon (NW Mediterranean), *J. Mar. Syst.* 60 (2006) 75–97.
- [210] D. Zuñiga, M.M. Flexas, A. Sanchez-Vidal, J. Coenjaerts, A. Calafat, G. Jorda, J. Garcia-Orellana, J. Puigdefabregas, M. Canals, M. Espino, et al., Particle fluxes dynamics in Blanes submarine canyon (Northwestern Mediterranean), *Prog. Oceanogr.* 82 (4) (2009) 239–251.
- [211] J.A. Salvadó, J.O. Grimalt, J.F. López, A. Palanques, S. Heussner, C. Pasqual, A. Sanchez-Vidal, M. Canals, Transfer of lipid molecules and polycyclic aromatic hydrocarbons to open marine waters by dense water cascading events, *Prog. Oceanogr.* 159 (2017) 178–194.
- [212] E. Lipiatou, J.C. Marty, A. Salot, Sediment trap fluxes of polycyclic aromatic hydrocarbons in the Mediterranean Sea, *Mar. Chem.* 44 (1993) 43–54.
- [213] C. Raoux, J.M. Bayona, J.C. Miquel, J.L. Teyssie, S.W. Fowler, J. Albaiges, Particulate fluxes of aliphatic and aromatic hydrocarbons in near-shore waters to the northwestern Mediterranean Sea, and the effect of continental runoff, *Estuar. Coast Shelf Sci.* 48 (1999) 605–616.
- [214] J. Dachs, J.M. Bayona, S.W. Fowler, J.-C. Miquel, J. Albaigés, Vertical fluxes of polycyclic aromatic hydrocarbons and organochlorine compounds in the western Alboran Sea (southwestern Mediterranean), *Mar. Chem.* 52 (1996) 75–86.
- [215] I. Bouloubassi, L. Mejanelle, R. Pete, J. Fillaux, A. Lorre, V. Point, PAH transport by sinking particles in the open Mediterranean Sea: a 1-year sediment trap study, *Mar. Pollut. Bull.* 52 (2006) 560–571.
- [216] M. Tsapakis, M. Apostolaki, S. Eisenreich, E.G. Stephanou, Atmospheric deposition and marine sedimentation fluxes of polycyclic aromatic hydrocarbons in the eastern Mediterranean basin, *Environ. Sci. Technol.* 40 (2006) 4922–4927.
- [217] J. Bonnin, S. Heussner, A. Calafat, J. Fabres, A. Palanques, X. Durrieu de Madron, M. Canals, P. Puig, J. Avril, N. Delsaut, Comparison of horizontal and downward particle fluxes across canyons of the Gulf of Lions (NW Mediterranean): meteorological and hydrodynamical forcing, *Cont. Shelf Res.* 28 (2008) 1957–1970.
- [218] A. Palanques, J. Guillen, P. Puig, X. Durrieu de Madron, Storm-driven shelf-to-canyon suspended sediment transport at the southwestern Gulf of Lions, *Cont. Shelf Res.* 28 (2008) 1947–1956.
- [219] S.M. Harmon, The toxicity of persistent organic pollutants to aquatic organisms, in: Y. Zeng (Ed.), Persistent Organic Pollutants (POPs): Analytical Techniques, Environmental Fate and Biological Effects, vol. 67, Comprehensive Analytical Chemistry, 2015, pp. 587–613.
- [220] X. Durrieu de Madron, C. Guieu, R. Sempéré, P. Conan, D. Cossa, F. D'Ortenzio, C. Estournel, F. Gazeau, C. Rabouille, L. Stemmann, S. Bonnet, F. Diaz, P. Koubbi, O. Radakovitch, M. Babin, M. Baklouti, C. Bancon-Montigny, S. Belviso, N. Bensoussan, B. Bonsang, I. Bouloubassi, C. Brunet, J.-F. Cadiou, F. Carlotti, M. Chami, S. Charmasson, B. Charrière, J. Dachs, D. Doxaran, J.-C. Dutay, F. Elbaz-Poulichet, M. Eléaume, F. Eyrolles, C. Fernandez, S. Fowler, P. Francour, J.C. Gaertner, R. Galzin, S. Gasparini, J.-F. Ghiglione, J.-L. Gonzalez, C. Goyet, L. Guidi, K. Guizien, L.-E. Heimbürger, S.H.M. Jacquet, W.H. Jeffrey, F. Joux, P. Le Hir, K. Leblanc, D. Lefèvre, C. Lejeune, R. Lemé, M.-D. Lojé-Pilot, M. Mallet, L. Méjanelle, F. Mélin, C. Mellon, B. Méricot, P.-L. Merle, C. Migon, W. L. Miller, L. Mortier, B. Mostajir, L. Mousseau, T. Moutin, J. Para, T. Pérez, A. Petrenko, J.-C. Poggiale, L. Prieur, M. Pujo-Pay, Pulido-Villena, P. Raimbault, A.P. Rees, C. Ridame, J.-F. Rontani, D. Ruiz Pino, M.A. Sicre, V. Taillandier, C. Tamburini, T. Tanaka, I. Taupier-Letage, M. Tedetti, P. Testor, H. Thébaud, B. Thouvenin, F. Touratier, J. Tronczynski, C. Ulsef, F. Van Wambeke, V. Vantrepotte, S. Vaz, R. Verney, Marine ecosystems' responses to climatic and anthropogenic forcings in the Mediterranean, *Prog. Oceanogr.* 91 (2011) 97–166.
- [221] M. Harmelin-Vivien, X. Bodiguel, S. Charmasson, V. Loizeau, C. Mellon-Duval, J. Tronczynski, D. Cossa, Differential biomagnification of PCB, PBDE, Hg and Radiocesium in the food web of the European hake from the NW Mediterranean, *Mar. Pollut. Bull.* 64 (5) (2012) 974–983.
- [222] M.J. Lawrence, H.I.J. Stemberger, A.J. Zolderdo, D.P. Struthers, S.J. Cooke, The effects of modern war and military activities on biodiversity and the environment, *Environ. Rev.* 23 (4) (2015) 443–460.
- [223] S. Guerzoni, R. Chester, F. Dulac, B. Herut, M.D. Loye-Pilot, C. Measures, C. Migon, E. Molinaroli, C. Moulin, P. Rossini, C. Saydam, A. Soudine, P. Ziveri, The role of atmospheric deposition in the biogeochemistry of the Mediterranean Sea, *Prog. Oceanogr.* 44 (1–3) (1999) 147–190.
- [224] N.H. Morley, J.D. Burton, S.P.C. Tankere, J.M. Martin, Distribution and behaviour of some dissolved trace metals in the western Mediterranean Sea, *Deep-Sea Res. Part II* 44 (3–4) (1997) 675–691.
- [225] EC, No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs, *Off. J. Eur. Union* (2006). L364/5-24.
- [226] EC, DIRECTIVE 2008/105/EC OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 16 December 2008 on environmental quality standards in the field of water policy, amending and subsequently repealing Council Directives 82/176/EEC, 83/513/EEC, 84/156/EEC, 84/491/EEC, 86/280/EEC and amending Directive 2000/60/EC of the European Parliament and of the Council, *Off. J. Eur. Union* (2008). L348/84-97.
- [227] EC, COMMISSION DECISION (EU) 2017/848 of 17 May 2017 laying down criteria and methodological standards on good environmental status of marine waters and specifications and standardised methods for monitoring and assessment, and repealing Decision 2010/477/EU, *Off. J. Eur. Union* (2017). L125/43-74.
- [228] A. Fliedner, H. Rüdell, B. Knopf, N. Lohmann, M. Paulus, M. Jud, U. Pirntke, J. Koschorreck, Assessment of seafood contamination under the marine strategy framework directive: contributions of the German environmental specimen bank, *Environ. Sci. Pollut. Control Ser.* 25 (27) (2018) 26939–26956.
- [229] M. Solé, C. Porte, J. Albaiges, Hydrocarbons, PCBs and DDT in the NW Mediterranean deep-sea fish *Mora moro*, *Deep Sea Res., Part I* 48 (2001) 495–513.
- [230] M. Trudel, J.B. Rasmussen, Predicting mercury concentration in fish using mass balance models, *Ecol. Appl.* 11 (2001) 517–529.
- [231] B.S. Galil, A. Golik, M. Türkay, Litter at the bottom of the sea: a sea bed survey in the Eastern Mediterranean, *Mar. Pollut. Bull.* 30 (1) (1995) 22–24, [https://doi.org/10.1016/0025-326X\(94\)00103-G](https://doi.org/10.1016/0025-326X(94)00103-G).
- [232] M.M. Storelli, S. Losada, G.O. Marcotrigiano, L. Roosens, G. Barone, H. Neels, A. Covaci, Polychlorinated biphenyl and organochlorine pesticide contamination signatures in deep-sea fish from the Mediterranean Sea, *Environ. Res.* 109 (2009) 851–856.
- [233] S. Damiano, P. Papetti, P. Menesatti, Accumulation of heavy metals to assess the health status of swordfish in a comparative analysis of Mediterranean and Atlantic areas, *Mar. Pollut. Bull.* 62 (2011) 1920–1925.
- [234] S. Koenig, P. Fernandez, M. Sole, Differences in cytochrome P450 enzyme activities between fish and crustacea: relationship with the bioaccumulation patterns of polychlorobiphenyls (PCBs), *Aquat. Toxicol.* 108 (2012) 11–17.
- [235] S. Koenig, M. Sole, C. Fernandez-Gomez, S. Diez, New insights into mercury bioaccumulation in deep-sea organisms from the NW Mediterranean and their human health implications, *Sci. Total Environ.* 442 (2013) 329–335.
- [236] D. Cossa, M. Coquery, The Mediterranean mercury anomaly, a geochemical or a biological issue, in: A. Salot (Ed.), The Mediterranean Sea - the Handbook of Environmental Chemistry, Springer-Verlag, Berlin, 2005, pp. 177–208, n° 5, Part K.
- [237] D. Cossa, M. Harmelin-Vivien, C. Mellon-Duval, V. Loizeau, B. Averty, S. Crochet, L. Chou, J.F. Cadiou, Influences of bioavailability, trophic position, and growth on methylmercury in hakes (*Merluccius merluccius*) from northwestern mediterranean and northeastern atlantic, *Environ. Sci. Technol.* 46 (2012) 4885–4893.
- [238] T. Chouvelon, P. Cresson, M. Bouchoucha, C. Brach-Papa, P. Bustamante, S. Crochet, F. Marco-Miralles, B. Thomas, J. Knoery, Oligotrophy as a major driver of mercury bioaccumulation in medium-to high-trophic level consumers: a

- marine ecosystem-comparative study, *Environ. Pollut.* 233 (Supplement C) (2018) 844–854.
- [239] J.C. Drazen, R.L. Haedrich, A continuum of life histories in deep-sea demersal fishes, *Deep-Sea Res.*, Part I 61 (2012) 34–42.
- [240] S. Koenig, P. Fernandez, J.B. Company, D. Huertas, M. Sole, Are deep-sea organisms dwelling within a submarine canyon more at risk from anthropogenic contamination than those from the adjacent open slope? A case study of Blanes canyon (NW Mediterranean), *Prog. Oceanogr.* 118 (2013) 249–259.
- [241] S. Koenig, D. Huertas, P. Fernández, Legacy and emergent persistent organic pollutants (POPs) in NW Mediterranean deep-sea organisms, *Sci. Total Environ.* 443 (2013) 358–366.
- [242] G. Rotllant, E. Abad, F. Sarda, M. Abalos, J.B. Company, J. Rivera, Dioxin compounds in the deep-sea rose shrimp *Aristeus antennatus* (Risso, 1816) throughout the Mediterranean Sea, *Deep-Sea Res.*, Part I 53 (2006) 1895–1906.
- [243] M. Carbery, W. O'Connor, T. Palanisami, Trophic transfer of microplastics and mixed contaminants in the marine food web and implications for human health, *Environ. Int.* 15 (2018) 400–409.
- [244] M.M. Storelli, A. Storelli, R. D'Addabbo, G. Barone, G.O. Marcotrigiano, Polychlorinated biphenyl residues in deep-sea fish from Mediterranean Sea, *Environ. Int.* 30 (2004) 343–349.
- [245] M.M. Storelli, V.G. Perrone, G.O. Marcotrigiano, Organochlorine contamination (PCBs and DDTs) in deep-sea fish from the Mediterranean Seasea, *Mar. Pollut. Bull.* 54 (2007) 1968–1971.
- [246] C. Porte, E. Escartin, L.M. Garcia, M. Solé, J. Albaigés, Xenobiotic metabolizing enzymes and antioxidant defences in deep-sea fish: relationship with contaminant body burden, *Mar. Ecol. Prog. Ser.* 192 (2000) 259–266.
- [247] F. Galgani, J.P. Leaute, P. Moguedal, A. Souplet, Y. Verin, A. Carpentier, H. Goraguier, D. Latrouite, B. Andral, Y. Cadiou, J.C. Mahe, J.C. Poulard, P. Nerisson, Litter on the sea floor along European coasts, *Mar. Pollut. Bull.* 40 (6) (2000) 516–527.
- [248] E. Ramirez-Llodra, B. De Mol, J.B. Company, M. Coll, F. Sardà, Effects of natural and anthropogenic processes in the distribution of marine litter in the deep Mediterranean Sea, *Prog. Oceanogr.* 118 (2013) 273–287.
- [249] C. Pham, E. Ramirez-Llodra, H.S. Claudia, T. Amaro, M. Bergmann, M. Canals, J. Company, J. Davies, G. Duineveld, F. Galgani, K. Howell, V.A. Huvnenne, E. Isidro, D. Jones, G. Lastras, T. Morato, J. Gomes-Pereira, A. Purser, H. Stewart, I. Tojeira, X. Tubau, D. Van Rooij, P. Tyler, Marine litter distribution and density in European seas, from the shelves to deep basins, *PLoS One* 9 (4) (2014), e95839.
- [250] C. Ioakeimidis, C. Zeri, E. Kaberi, M. Galatchi, K. Antoniadis, N. Streftaris, F. Galgani, E. Papatheodorou, G. Papatheodorou, A comparative study of marine litter on the seafloor of coastal areas in the Eastern Mediterranean and Black Seas, *Mar. Pollut. Bull.* 89 (2014), 296–30.
- [251] M. Angiolillo, B. Lorenzo, A. Farcomeni, M. Bo, G. Bavestrello, G. Santangelo, A. Cau, V. Mastascusa, F. Sacco, S. Canese, Distribution and assessment of marine debris in the deep Tyrrhenian Sea (NW Mediterranean sea, Italy), *Mar. Pollut. Bull.* 92 (2015) 149–159.
- [252] M. Pierdomenico, D. Casalbore, F.L. Chiocci, Massive benthic litter funnelled to deep sea by flash-flood generated hypervelocity flows, *Sci. Rep.* 9 (2019) 5330.
- [253] J.P.W. Desforges, M. Galbraith, N. Dangerfield, P.S. Ross, Widespread distribution of microplastics in subsurface seawater in the NE Pacific Ocean, *Mar. Pollut. Bull.* 79 (2014) 94–99.
- [254] L.C. Woodall, A. Sanchez-Vidal, M. Canals, G.L.J. Paterson, R. Coppock, V. Sleight, A. Calafat, A.D. Rogers, B.E. Narayanaswamy, R.C. Thompson, The Deep Sea Is a Major Sink for Microplastic Debris, vol. 1, Royal Society Open Science, 2014, p. 140317.
- [255] L. Van Cauwenbergh, A. Vanreusel, J.C. Maes, Microplastic pollution in deep sea sediments, *Environ. Sci. Pollut. Control Ser.* 182 (2013) 495–499.
- [256] A. Sanchez-Vidal, R.C. Thompson, M. Canals, W.P. de Haan, The imprint of microfibrils in southern European deep seas, *PLoS One* 13 (11) (2018), e0207033.
- [257] M. Gregory, Environmental implications of plastic debris in marine settings entanglement, ingestion, smothering, hangers-on, hitch-hiking and alien invasions, *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 364 (1526) (2009) 2013–2025.
- [258] F. Murray, P. Cowie, Plastic contamination in the decapod crustacean *Nephrops norvegicus* (Linnaeus, 1758), *Mar. Pollut. Bull.* 62 (2011) 1207–1217.
- [259] G.A. Anastasopoulou, C. Mytilineou, C. Smith, K. Papadopoulou, Plastic debris ingested by deep-water fish of the Ionian Sea (eastern mediterranean), *Deep Sea Res.* 1 74 (2012) 11–13.
- [260] M.L. Taylor, C. Gwinnett, L.F. Robinson, L.C. Woodall, Plastic microfibre ingestion by deep-sea organisms, *Sci. Rep.* 6 (2016) 33997.
- [261] S. Newman, E. Watkins, A. Farmer, P. Ten Brink, J.P. Schweitzer, The Economics of marine litter, in: M. Bergmann, L. Gutow, M. Klages (Eds.), *Marine Anthropogenic Litter*, Springer, Cham, 2015.
- [262] J. Oehlmann, U. Schulte-Oehlmann, W. Kloas, O. Jagnytsh, I. Lutz, K. Kusk, L. Wollenberger, E. Santos, G. Paull, K. Van Look, C. Tyler, A critical analysis of the biological impacts of plasticizers on wildlife, *Phil. Trans. R. Soc. Part B* 364 (2009) 2047–2062.
- [263] D.P. Nowacek, C.W. Clark, D. Mann, P.J. Miller, H.C. Rosenbaum, J.S. Golden, M. Jasný, J. Kraska, B.L. Southall, Marine seismic surveys and ocean noise: time for coordinated and prudent planning, *Front. Ecol. Environ.* 13 (2015) 378–386.
- [264] R. Rountree, J. Aguzzi, S. Marini, E. Fanelli, F.C. De Leo, J. Del Rio, F. Juanes, Towards an optimal design for ecosystem-level ocean observatories, *Adv. Mar. Biol.*, *Ann. Rev.* (2019) in press.
- [265] J.C. Montgomery, C.A. Radford, Marine bioacoustics, *Curr. Biol.* 27 (2017) R502–R507.
- [266] S.D. Simpson, M. Meekan, J. Montgomery, R. McCauley, A. Jeffs, Homeward sound, *Science* 308 (5719) (2005), <https://doi.org/10.1126/science.1107406>, 221–221.
- [267] D. Wartzk, D.R. Ketten, Marine mammal sensory systems, in: J. Reynolds, S. Rommel (Eds.), *Biology of Marine Mammals*, Smithsonian Institution Press, Washington, DC, 1999, ISBN 1560983752, pp. 117–175.
- [268] P.L. Edds-Walton, Acoustic communication signals of mysticete whales, *Bioacoustics* 8 (1997) 47–60.
- [269] W.W.L. Au, *The Sonar of Dolphins*, Springer-Verlag, New York, 1993.
- [270] G.M. Wenz, Acoustic ambient noise in the ocean: spectra and sources, *J. Acoust. Soc. Am.* 34 (1962) 1936–1956.
- [271] J.A. Hildebrand, Anthropogenic and natural sources of ambient noise in the ocean, *Mar. Ecol. Prog. Ser.* 395 (2009) 5–20, <https://doi.org/10.3354/meps08353>.
- [272] A. Maglio, G. Pavan, M. Castellote, S. Frey, Overview of the noise hotspots in the ACCOBAMS area – Part I, Mediterranean Sea, in: *Agreement on the Conservation of Cetaceans in the Black Sea, Mediterranean Sea and Contiguous Area*, 2016, p. 44.
- [273] A.N. Popper, A.D. Hawkins, R.R. Fay, D.A. Mann, S. Bartol, T.J. Carlson, S. Coombs, W.T. Ellison, R.L. Gentry, M.B. Halvorsen, S. Lokkeborg, P. Rogers, B. L. Southall, D.G. Zeddis, W.N. Tavolga, Sound Exposure Guidelines for Fishes and Sea Turtles: A Technical Report Prepared by ANSI-Accredited Standards Committee S3/SC1 and Registered with ANSI, Springer Briefs in Oceanography, 2014. <http://www.springer.com/gp/book/9783319066585>.
- [274] C. Peng, X.G. Zhao, G.X. Liu, Noise in the sea and its impacts on marine organisms, *Int. J. Environ. Res. Public Health* 12 (2015) 12304–12323.
- [275] A.D. Foote, R.W. Osborne, A.R. Hoelzel, Environment - whale-call response to masking boat noise, *Nature* 428 (6986) (2004), <https://doi.org/10.1038/428910a>, 910–910.
- [276] M. André, M. Solé, M. Lenoir, M. Durfurt, C. Quero, A. Mas, A. Lombarte, M. van der Schaer, M. Lopez-Bejar, M. Morell, et al., Low-frequency sounds induce acoustic trauma in cephalopods, *Front. Ecol. Environ.* 9 (2011) 489–493.
- [277] A. Azzellino, C. Lanfredi, A. D'Amico, G. Pavan, M. Podestà, J. Haun, Risk mapping for sensitive species to underwater anthropogenic sound emissions: model development and validation in two Mediterranean areas, *Mar. Pollut. Bull.* 63 (2011) 56–70.
- [278] A.G. Carroll, R. Przeslawski, A. Duncan, M. Gunning, B. Bruce, A critical review of the potential impacts of marine seismic surveys on fish & invertebrates, *Mar. Pollut. Bull.* 114 (2017) 9–24.
- [279] S.E. Nelms, W.E.D. Piniak, C.R. Weir, B.J. Godley, Seismic surveys and marine turtles: an underestimated global threat? *Biol. Conserv.* 193 (2016) 49–65.
- [280] S.L. DeRuiter, K. Larbi Doukara, Loggerhead turtles dive in response to airgun sound exposure, *Endanger. Species Res.* 16 (2012) 55–63.
- [281] L. Weiglart, The Impact of Ocean Noise Pollution on Fish and Invertebrates, Report for OceanCare, OCEANCARE & DALHOUSIE UNIVERSITY, 2017, https://www.oceancare.org/wp-content/uploads/2017/10/noise_and_fish_review_paper-20171016.pdf.
- [282] R. McCauley, R.D. Day, K.M. Swadling, Q.P. Fitzgibbon, R.A. Watson, Widely used marine seismic survey air gun operations negatively impact zooplankton, *Nat. Ecol. Evol.* 1 (2017) 1–8, <https://doi.org/10.1038/s41559-017-0195>.
- [283] F. Caruso, V. Sciacca, G. Bellia, E. De Domenico, G. Larosa, E. Papale, C. Pellegrino, S. Pulvirenti, G. Riccobene, F. Simeone, KM3NET Consortium, Size distribution of sperm whales acoustically identified during long term deep-sea monitoring in the Ionian Sea, *PLoS One* 10 (2015) 12.
- [284] P. Favali, F. Chierici, G. Marinaro, G. Giovanetti, A. Azzarone, L. Beranzoli, KMNET consortium, NEMO-SN1 abyssal cabled observatory in the Western Ionian Sea, *IEEE J. Ocean. Eng.* 38 (2013) 358–374.
- [285] M. André, A. Caballé, M. Van Der Schaer, A. Solsona, L. Houégnigan, S. Zaugg, ANTARES consortium, Sperm whale long-range echolocation sounds revealed by ANTARES, a deep-sea neutrino telescope, *Sci. Rep.* 7 (2017) 45517.
- [286] NRC-National Research Council (US), Committee on Potential Impacts of Ambient Noise in the Ocean on Marine Mammals, National Academies Press (US), Washington (DC), 2003.
- [287] S. Bianchelli, E. Buschi, R. Danovaro, A. Pusceddu, Nematode biodiversity and benthic trophic state are simple tools for the assessment of the environmental quality in coastal marine ecosystems, *Ecol. Ind* 95 (2018) 270–285.
- [288] M. Taviani, L. Angeletti, L. Beuck, E. Campiani, S. Canese, F. Fogliani, A. Freiwald, P. Montagna, F. Trincardi, On and off the beaten track: Megafaunal sessile life and Adriatic cascading processes, *Mar. Geol.* 369 (2015) 273–274.
- [289] A. Izquierdo-Muñoz, D. Izquierdo-Gomez, et al., First record of Lagocephalus scleratus (Gmelin, 1789) (Actinopterygii, Tetraodontidae) on the Mediterranean Spanish coast, in: S. Katsanevakis (Ed.), *New Mediterranean Biodiversity Records*, 2014, pp. 686–687. *Med. Mar. Sci.* 15(3), 675–695.
- [290] EU, Regulation (EC) No 45/2001 of the European Parliament and of the Council of 18 December 2000 on the protection of individuals with regard to the processing of personal data by the Community institutions and bodies and on the free movement of such data, 2000.
- [291] S.E. Moffitt, R.A. Moffitt, W. Sauthoff, C.V. Davis, K. Hewett, T.M. Hill, Paleooceanographic insights on recent oxygen minimum zone expansion: lessons for modern oceanography, *PLoS One* 10 (1) (2015).
- [292] D.S.M. Billeit, B.J. Bett, W.D.K. Reid, B. Boorman, I.G. Priede, Long-term change in the abyssal NE Atlantic: The 'Amperima Event' revisited, *Deep Sea Res.* II 57 (15) (2010) 1406–1417.
- [293] A. Pusceddu, S. Frascchetti, M. Scopa, L. Rizzo, R. Danovaro, Meiofauna communities, nematode diversity and C degradation rates in seagrass (*Posidonia*

- oceanica L.) and unvegetated sediments invaded by the algae *Caulerpa cylindracea* (Sonder), *Mar. Env. Res.* 119 (2016) 88–99.
- [294] V. Lauria, G. Garofalo, M. Gristina, F. Fiorentino, Contrasting habitat selection amongst cephalopods in the Mediterranean Sea: when the environment makes the difference, *Mar. Env. Res.* 119 (2016) 252–266.
- [295] E. Ramirez-Llodra, H.C. Trannum, A. Evensen, L.A. Levin, M. Andersson, T. E. Finne, A. Hilario, B. Flem, G. Christensen, M. Schaanning, et al., Submarine and deep-sea mine tailing placements: A review of current practices, environmental issues, natural analogs and knowledge gaps in Norway and internationally, *Mar. Poll. Bull.* 97 (1-2) (2015) 13–35.
- [296] K.O. Buesseler, The decoupling of production and particulate export in the surface ocean, *Global Biogeochem. Cycles* 12 (2) (1998) 297–310.
- [297] UNEP, *Marine Litter Assessment in the Mediterranean*, UNEP/MAP, Athens, 2015, p. 45, 2015.